



- 540 -

Step D

The compound produced in Step C was subjected to the conditions described in Tetrahedron Letters, Vol. 25, 1984, 839-842 to reduce the nitro group. The reduction was performed on 2 g of nitro compound to give 1 g of product. MS and H-NMR were consistent with the proposed structure.

15 Step E

.TFA

The compound produced in Step D was guanidated according to the method in Example M on a 1 g scale and purified by reverse phase chromatography (water/acetonitrile) to result in 110 mg of the title compound as a white solid. MS and H-NMR were consistent with the proposed structure.

30 Step F

The compound produced in Step E (100 mg) was dissolved in water/acetonitrile (1:1), followed by the addition of lithium hydroxide (100 mg). The reaction was allowed to stir at 25°C, and monitored by HPLC. After



- 541 -

complete hydrolysis (1-2 hours) trifluoroacetic acid was added until pH = 2. The product was purified by reverse phase chromatography (water/acetonitrile) to result in 110 mg of the title compound as a white solid. MS and H-NMR  
5 were consistent with the proposed structure.

- 542 -


Example 290

Preparation of  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid, trifluoroacetate salt


Step A

N,N-disuccinimidyl carbonate (DSC) (65 g) was added to methyl hydrogen 5-nitroisophthalate (5 g) in dry dimethylformamide (10 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes tert-butyl  $\beta$ -glycine (2.6 g) was added followed by addition of NMM (2.0 mL). After complete reaction (4 hours) the product was isolated by extraction into ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$  to result in a white solid (5.1 g). MS and H-NMR were consistent with the proposed structure.

- 543 -

Step B

The compound produced in Step A (5 g) was dissolved in dioxane (50 mL). To this mixture dry HCl (20 mL, 4N) was added. The mixture was stirred for 12 hours. The solvent was removed under reduced pressure followed by addition of ether and removal of the solvent under reduced pressure. The solid was filtered to result in a white solid (4 g) and dried in a vacuum oven. MS and H-NMR were consistent with the proposed structure.

Step C


N,N'-disuccinimidyl carbonate (DSC) (2 g) was added to the compound produced in Step B (2 g) in dry dimethylformamide (10 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes DL-ethyl 3-amino-3-(3,5-dichlorophenyl)propionate hydrochloride (1.6 g) was added followed by addition of NMM (2.0 mL). After complete reaction (4 hours) the product was isolated by extraction into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in an oil (3 g). MS and H-NMR were consistent with the proposed structure.



- 544 -

Step D

5




The compound produced in Step C was subjected to the conditions described in Tetrahedron Letters, Vol. 25, 1984, 839-842 to reduce the nitro group. The reduction was performed on 2g of nitro compound to give 1 g of product.

Step E

15

20




The compound produced in Step D was guanidated according to the method in Example M on a 1 g scale and purified by reverse phase chromatography (water/acetonitrile) to result in 110 mg of the title compound as a white solid. MS and H-NMR were consistent with the proposed structure.

- 545 -

Step F

5





10       The compound produced in Step E (100 mg) was dissolved in water/acetonitrile (1:1), followed by the addition of lithium hydroxide (100 mg). The reaction was allowed to stir at 25°C, and monitored by HPLC. After complete hydrolysis (1-2 hours) trifluoroacetic acid was  
15       added until pH = 2. The product was purified by reverse phase chromatography (water/acetonitrile) to result in 25 mg of the title compound as a white solid. MS and H-NMR were consistent with the proposed structure.

20

- 546 -

Example 291

Preparation of  $\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzene-  
propanoic acid, bis(trifluoroacetate) salt


Step A

Glycine (20 g, 266 mmol) was added to water (200 mL), followed by addition of potassium hydroxide (20 g, 357 mmol) and cooled to 0°C in an ice bath. To this mixture 3,5-dinitrobenzoyl chloride (20 g, 108 mmol) was added in acetonitrile (20 mL) drop-wise over a 10 minute period. After complete reaction (3-4 hours) concentrated hydrochloric acid was added until pH=1. The product was filtered, washed with water and air dried (20 g). MS and H-NMR were consistent with the proposed structure.

- 547 -

Step B


5



10 N,N'-disuccinimidyl carbonate (DSC) (1.2 g) was added to the compound produced in Step A (2 g) in dry dimethylformamide (10 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes, DL ethyl 3-amino-3-(3,5 dichlorophenyl) propionate hydrochloride (1.2 g) was added followed by addition of NMM (2.0 mL). After complete reaction  
15 (4 hours) the product was isolated by extraction into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in a yellow oil (2.1 g). MS and H-NMR were consistent with the proposed structure.

20 Step C

25




30 The compound produced in Step B was subjected to the conditions described in Tetrahedron Letters, Vol. 25, 1984, 839-842 to reduce of the nitro group. The reduction was performed on 2.5 g of nitro compound to give 2.1 g of the 3,5-dianilino derivative. MS and H-NMR were consistent with the proposed structure.

- 548 -

Step D

5



10        The compound produced in Step C was guanidated  
according to the method in Example M on a 2 g scale (using  
4 g of the guanidating agent) and purified by reverse  
phase chromatography (water/acetonitrile) to result in  
800 mg of a white solid. MS and H-NMR were consistent  
15        with the proposed structure.

Step E

20        The compound produced in Step D (500 mg) was  
dissolved in water/acetonitrile (1:1), followed by  
addition of lithium hydroxide (100 mg). The reaction was  
allowed to stir at 25°C, and monitored by HPLC. After  
complete hydrolysis (1-2 hours) trifluoroacetic acid was  
added until pH = 2. The product was purified by reverse  
phase chromatography (water/acetonitrile) to result in  
25        450 mg of the title compound as a white solid. MS and  
H-NMR were consistent with the proposed structure.

- 549 -

Example 292


Preparation of  $\beta$ -[[2-[[[3-[(amin iminomethyl)amino]-5-(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoic acid, trifluoroacetate salt

5

10

15


20

Step A

A mixture of 5-amino-3-nitro benzoic acid (Lancaster) (3 g) and trifluoroacetic anhydride (Sigma) (20 mL) in methylene chloride was stirred for 2 days at 25°C. The solvent was removed under reduced pressure to leave an oil. To the oil was added water (50 mL) and the product filtered (4.5 g). The product was dried in an oven at 70°C for 16 hours. MS and H-NMR were consistent with the proposed structure.


30

- 550 -

Step B

5  
10       N,N'-disuccinimidyl carbonate (DSC) (3 g) was added to the compound produced in Step A (2.7 g) in dry dimethylformamide (4 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes, tert-butyl glycine hydrochloride (2.7 g) was  
15 added followed by addition of NMM (2.0 mL). After complete reaction (4 hours) the product was isolated by extraction into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in an yellow oil (3.3 g). MS and H-NMR were consistent with the proposed structure.

20

Step C


25

30       The compound produced in Step B (3 g) was dissolved in methylene chloride (50 mL). To this mixture TFA (20 mL) was added. The mixture was stirred for 12 hours. The solvent was removed under reduced pressure followed by addition of ether. The solid (2.7 g) was filtered and

- 551 -


dried in a vacuum oven for 1-2 hours. MS and H-NMR were consistent with the proposed structure.

Step D



10 N,N'-disuccinimidyl carbonate (DSC) (1.5 g) was added to the product of Step C (1.2 g) in dry dimethylformamide (10 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes DL ethyl 3-amino-3-(3,5  
15 dichlorophenyl) propionate hydrochloride (1.7 g) was added followed by addition of NMM (2.0 mL). After complete reaction (4 hours) the product was isolated by extraction into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in an yellow oil (2.1 g). MS and H-NMR were consistent with the  
20 proposed structure.

Step E



25  
30 The compound produced in Step D was subjected to the conditions described in Tetrahedron Letters, Vol. 25, 1984, 839-842 to reduce the nitro group. The reduction was performed on 1.8 g of nitro compound to give 1.8 g of




- 552 -

the 3-anilino derivative. MS and H-NMR were consistent with the proposed structure.

Step F

5

10



15 The compound produced in Step E was guanidated according to the method in Example M on a 1.5 g scale (using 3 g of the guanidating agent) and purified by reverse phase chromatography (water/acetonitrile) to result in 750 mg of the above compound as a white solid.

20 MS and H-NMR were consistent with the proposed structure.

Step G

The compound produced in Step F (500 mg) was dissolved in water/acetonitrile (1:1), followed by the



25 addition of lithium hydroxide (100 mg). The reaction was allowed to stir at 25°C, and monitored by HPLC. After complete hydrolysis (1-2 hours) trifluoroacetic acid was added until pH = 2. The product was purified by reverse phase chromatography (water/acetonitrile) to result in 300

30 mg of the title compound as a white solid. MS and H-NMR were consistent with the proposed structure.

- 553 -

Example 293

Preparation of  $\beta$ -[[2-[[[3-(acetylamino)-5-  
5 [(aminoiminomethyl)amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzenepropanoic  
acid, trifluoroacetate salt


Step A

A mixture of 5-amino-3-nitro benzoic acid (Lancaster)  
25 (5 g) and acetic anhydride (Sigma) (10 mL) in methylene  
chloride was stirred for 2 days at 25°C. The solvent was  
removed under reduced pressure to leave an oil. To the  
oil was added water (50 mL) and the product filtered  
(4.5 g). The product was dried in an oven at 70°C for 16  
30 hours. MS and H-NMR were consistent with the proposed  
structure.

- 554 -

Step B


5



10       N,N'-disuccinimidyl carbonate (DSC) (3 g) was added  
to the compound produced in Step A (3 g) in dry  
dimethylformamide (6 mL) followed by addition of  
dimethylaminopyridine (200 mg). After a period of 20  
minutes tert-butyl glycine hydrochloride (2.1 g) was added  
followed by addition of NMM (2.0 mL). After complete  
15       reaction (4 hours) the product was isolated by extraction  
into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in an  
yellow oil (3.3 g). MS and H-NMR were consistent with the  
proposed structure.

20   Step C

25




The compound produced in Step B (3 g) was dissolved  
in methylene chloride (10 mL). To this mixture TFA (10  
30       mL) was added. The mixture was stirred for 12 hours. The  
solvent was removed under reduced pressure followed by  
addition of addition of ether. The solid (3 g) was  
filtered and dried in a vacuum oven for 1-2 hours. MS and  
H-NMR were consistent with the proposed structure.

- 555 -

Step D

5




10 N,N'-disuccinimidyl carbonate (DSC) (1.5 g) was added to the product from Step C (1.2 g) in dry dimethylformamide (10 mL) followed by addition of dimethylaminopyridine (200 mg). After a period of 20 minutes DL ethyl 3-amino-3-(3,5 dichlorophenyl) propionate hydrochloride (1.7 g) was added followed by addition of NMM (2.0 mL). After complete reaction (4 hours) the product was isolated by extraction into ethyl acetate and dried over Na<sub>2</sub>SO<sub>4</sub> to result in a tan solid (2 g). MS and H-NMR were consistent with the proposed structure.

15

Step E

20




25

30 The compound produced in Step D was subjected to the conditions described in Tetrahedron Letters, Vol. 25, 1984, 839-842 to reduce the nitro group. The reduction was performed on 1.5 g of nitro compound to give 1.5 g of the 3-anilino derivative. MS and H-NMR were consistent with the proposed structure.

- 556 -

Step F

5




10

The compound produced in Step E was guanidated according to the method in Example M on a 1.4 g scale (using 2 g of the guanidating agent) and purified by reverse phase chromatography (water/acetonitrile) to result in 750 mg of the above compound as a white solid. MS and H-NMR were consistent with the proposed structure.

Step G

The compound produced in Step F (300 mg) was dissolved in water/acetonitrile (1:1), followed by the addition of lithium hydroxide (100 mg). The reaction was allowed to stir at 25°C, and monitored by HPLC. After complete hydrolysis (1-2 hours) trifluoroacetic acid was added until pH = 2. The product was purified by reverse phase chromatography (water/acetonitrile) to result in 200 mg of the title compound as a white solid. MS and H-NMR were consistent with the proposed structure.

- 557 -


Examples 294-296Step A

To a 2L-3-neck round bottom flask equipped with  
10 mechanical stirrer was added  $\beta$ -amino-3,5-dichloro-  
benzenepropanoic acid (52.78 g, 0.2255 mol). The  $\beta$ -amino-  
3,5-dichlorobenzenepropanoic acid was dissolved in 900 mL  
of acetone and 300 mL of water and sodium carbonate was  
added (3.0 eq., 71.70 g, 0.6765 mol). The pH = 10. The  
15 Fmoc succinimidyl carbonate (Sigma Chemical Co., 1.0 eq.,  
76.06 g, 0.2255 mol) was dissolved in 600 mL of acetone  
and added slowly to the basic aqueous solution via  
addition funnel over 45 minutes. The reaction was stirred  
for 16 hours at room temperature. HPLC analysis (Waters,  
20 C18, reverse phase, 25 cm column, 50-90% acetonitrile in  
water over 30 minutes) indicated that the  $\beta$ -amino-3,5-  
dichlorobenzenepropanoic acid was consumed. The acetone  
was removed from the reaction mixture *in vacuo*. The basic  
aqueous phase was acidified to pH = 3 using 3.0 N  
25 hydrochloric acid. In a 2L separatory funnel the acid  
layer was washed with 1L of ethyl acetate, the water layer  
was removed and the organic layer was washed (2 x 250 mL  
water, 2 x 250 mL saturated sodium chloride). The organic  
layer was dried (magnesium sulfate), filtered and  
30 concentrated *in vacuo* to 300 mL. Petroleum ether was  
added (300 mL) and a white flocculent solid precipitated.  
After 24 hours of air drying, isolated 38.49 g as a first  
crop (38% yield). The mother liquor was saved for future  
use. NMR (DMSO): 2.62-2.72 (m, 2H), 4.15-4.32 (m, 1H),  
35 7.21-7.40 (m, 5H), 7.45 (s, 1H), 7.60-7.70 (m, 2H), 7.85

- 558 -

(d,  $j=7$  Hz, 2H), 7.99 (d,  $j=7$  Hz, 1H). MS (FAB)  $m/e$  (relative intensity): 456.2 (20), 179 (100).

### Step B




Wang resin (25.0 g, 28.0 mmol) was placed in a 1L 3-neck round bottom flask fitted with an overhead stirrer and nitrogen inlet. The resin was swelled with 250 mL of methylene chloride for 15 minutes then drained. The FMOC protected amino acid produced in Step A (25.66 g, 56.0 mmol) was activated in a separate 500 mL round bottom flask by dissolving in methylene chloride/dimethylformamide (4:1, 125 mL) and adding diisopropylcarbodiimide (DIC, 8.77 mL, 56.0 mmol) via syringe, followed by addition of dimethylaminopyridine (DMAP, 0.342 g, 2.8 mmol). The solution was stirred at 25°C for 15 minutes, then added to the preswelled Wang resin. The slurry was stirred for 2 hours at 25°C. The reaction was drained and washed with methanol (3 x 250 mL), methylene chloride (3 x 250 mL) and diethyl ether (3 x 250 mL). The resin was then swelled in 250 mL of methylene chloride and drained. The activated product of Step A (12.83 g, 28.0 mmol, DIC, 4.36 mL, 28.0 mmol, DMAP, 0.170 g, 1.4 mmol in 100 mL methylene chloride/dimethylformamide 4:1) was added to the swelled resin. The slurry was stirred at 25°C for 1 hour. The resin was drained and washed as before. Elemental analysis calculated for resin bound material:

Calculated: C, 81.31; H, 6.30; N, 1.05; Cl, 5.33.

Found: C, 79.03; H, 6.37; N, 1.16; Cl, 5.74.

- 559 -

Step C

5

10 The product of Step B (28.0 mmol) was preswelled in a 1L 3-neck round bottom flask equipped with overhead stirrer and nitrogen inlet using 250 mL of methylene chloride for 15 minutes. The solvent was drained and a 20% piperidine/dimethylformamide solution (125 mL) was added and the slurry was stirred at 25°C for 2 hours. The

15 resin was drained and washed with dimethylformamide (3 x 100 mL), methanol (3 x 100 mL) methylene chloride (3 x 100 mL) and diethyl ether (3 x 100 mL). The resin was dried using house vacuum for 1 hour. An activated solution of FMOC-Glycine (20.81 g, 70.0 mmol, DIC, 10.95 mL, 70.0

20 mmol, DMAP, 0.85 g, 7.0 mmol. In 150 mL methylene chloride/dimethylformamide, 4:1) was added to the preswelled resin via syringe and stirred at 25°C for 2 hours. The resin was drained and washed (methylene chloride, methanol and diethyl ether, each 3 x 100 mL).

25 The resin was preswelled with 250 mL of methylene chloride for 15 minutes, drained and a solution of activated FMOC-Glycine (10.45 g, 35.0 mmol, DIC, 5.42 mL, 35.0 mmol, DMAP, 0.42 g, 3.5 mmol in 100 mL methylene chloride/dimethyl formamide 4:1) was added to the swelled


30 resin via syringe. The slurry was stirred at 25°C for 1 hour. The resin was drained and washed (methylene chloride, methanol, diethyl ether, 3 x 100 mL each). The resin was vacuum dried for 1 hour. The Kaiser test (Kaiser, E., Color Test for Detection of Free Terminal



- 560 -


Amino Groups in the Solid-Phase Synthesis of P ptid s.  
Anal. Biochem. 1970, 34, 595-598) indicated c upling was complete.

5 Step D



In a 500 mL bottom flask equipped with magnetic stirrer, 3-amino-benzoic acid (Aldrich, 10.0 g, 50.8 mmol) was dissolved in 50 mL of dioxane and 133 mL of 10% sodium carbonate. The stirred solution was cooled to 0°C  
10 (ice/water) and a solution of fluorenylmethyl chloroformate (13.78 g, 53.3 mmol, in 50 mL dioxane) was added dropwise over 15 minutes. The reaction was warmed to 25°C overnight. HPLC analysis (as described earlier) indicated that the starting material was consumed. 500 mL  
15 of water was added to the reaction mixture and a white precipitate formed immediately. The solid was collected, washed with 10% citric acid and dried under vacuum. Isolated 15.23 g, (83.4% yield) of a white flocculent solid. NMR (DMSO): 4.18-4.25 (m, 3H), 7.25-7.41 (m, 6H),  
20 7.62-7.72 (m, 3H), 7.89-7.90 (m, 3H). MS(FAB): product ion  $\text{M}+\text{H}$  observed at  $m/z$  360.

- 561 -

Step E

20.0 g of the product of Step C (22.4 mmol) was preswelled in 500 mL of methylene chloride for 30 minutes. The solvent was drained and 250 mL of 20% piperidine/dimethyl formamide was added and allowed to stir at 25°C for 40 minutes. The resin was drained and washed with dimethyl formamide, methanol, methylene chloride, and diethyl ether (each solvent, 3 x 150 mL). The Kaiser test indicated the deprotection was complete. The resin was dried using house vacuum for 45 minutes. The resin was then preswelled using 250 mL of methylene chloride, drained and the activated product of Step D (13.54 g, 35.5 mmol, DIC, 5.55 mL, 35.5 mmol, DMAP, 0.88 g, 7.2 mmol, in 100 mL methylene chloride/dimethyl formamide 4:1) was added to the preswelled resin. The reaction was stirred for 16 hours at 25°C. The resin was drained and washed as previously described. The Kaiser test indicated that the reaction was not complete. The coupling reaction was repeated, the resin was drained and washed. A repeat Kaiser test indicated that the coupling reaction was complete. A small portion of the resin was Fmoc deprotected (30 minutes with 20% piperidine/dimethyl formamide) then cleaved off resin (1 hour with 95% trifluoroacetic acid/water) for NMR analysis. NMR (DMSO): 2.68-2.78 (m, 2H), 3.88 (d,  $j=7$  Hz, 2H), 5.06-5.20 (m, 1H), 7.32-7.69 (m, 4H), 7.54 (t,  $j=8$  Hz, 1H), 7.76


- 562 -

(s, 1H), 7.83 (d, j=8 Hz, 1H), 8.57 (d, j=9 Hz, 1H), 8.87 (t, j=9 Hz, 1H).

Step F

5

10




The resin of Step E (2.0 g, 2.0 mmol) in a 100 mL round bottom flask, was preswelled with 20 mL of dimethyl formamide, drained, then treated with 20 mL of 20% piperidine/dimethyl formamide for 40 minutes at 25°C. The resin was filtered and washed with dimethyl formamide, methanol, methylene chloride and diethyl ether (3 x 10 mL, each). The Kaiser test was inconclusive, and the deprotection step and washings were repeated. The repeat Kaiser test was still inconclusive, and the material used as is. The 2.0 g of resin was split into two 1.0 g portions and placed into 2 dram glass vials. Dimethyl formamide (4.0 mL/vial) was added, followed by methyl isothiocyanate (1.4622 g, 20 mmol). The vials were tightly capped and heated to 80°C for 4 hours. The resin was filtered and washed with dimethyl formamide, methanol, methylene chloride, and diethyl ether (3 x 10 mL, each). The resin was dried in vacuo.

30

- 563 -

Step G

5




The resin product from Step F was transferred to a  
10 fritted, 100 mL reaction vessel. The resin was swelled  
with methylene chloride (3 x 10 mL) and drained. In a  
separate vial 2-chloro-1-methylpyridiniumiodide (Aldrich,  
0.405 g, 1.58 mmol) was dissolved in 5 mL of  
dimethylformamide/methylene chloride 4:1 and added to the  
15 preswelled resin, followed by triethylamine (0.441 mL,  
3.17 mmol). The reaction slurry was stirred for 8 hours  
at 25°C. The resin was drained, and washed with  
dimethylformamide and methylene chloride (3 x 10 mL  
each). The resin was dried *in vacuo*.

20

Step H

25




The resin product from Step G (0.666 g, 0.7 mmol) was  
30 transferred to a 15 mL fritted vessel and suspended in 3.5  
mL of dimethylformamide/methylene chloride (1:1).  
Methylamine (2.0 M in tetrahydrofuran, 4.4 mL, 8.8 mmol)  
was added to the resin slurry and allowed to stir at 25°C  
for 16 hours. The resin was drained, and washed with

- 564 -


dimethylformamide, methanol, methylene chloride and diethyl ether (3 x 10 mL each). The resin was dried in vacuo for 1 hour.

**5 Step I**



The resin product from Step G (0.666 g, 0.7 mmol) was transferred to a 15 mL fritted vessel and suspended in 3.5 mL dimethylformamide/methylene chloride (1:1). Ethylamine (2.0 M in tetrahydrofuran, 4.4 mL, 8.8 mmol) was added to the resin slurry and allowed to stir at 25°C for 16 hours. The resin was drained, and washed with dimethylformamide, methanol, methylene chloride and diethyl ether (3 x 10 mL each). The resin was dried in vacuo for 1 hour.

**Step J**




- 565 -

The resin product from St p G (0.666 g, 0.7 mmol) was transferred to a 15 mL fritted vessel and suspended in 3.5 mL dimethylformamide/methylene chloride (1:1). Isopropylamine (0.749 mL, 8.8 mmol) was added to the resin  
5 slurry and allowed to stir at 25°C for 16 hours. The resin was drained, and washed with dimethylformamide, methanol, methylene chloride and diethyl ether (3 x 10 mL each). The resin was dried *in vacuo* for 1 hour.

- 566 -

Example 294

Preparation of ( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-  
[[ (methylamino) (methylimino) methyl] amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
benzenepropanoic acid,  
trifluoroacetate salt




The resin product from Step H was treated with 2.5 mL of 95% trifluoroacetic acid/water for 1 hour at 25°C. The filtrate was collected. The resin was washed with 2 x 1 mL of 50% trifluoroacetic acid/methylene chloride and the filtrate collected. The resin was washed once more with 1 mL of methylene chloride. All filtrates were combined into a tared 2 dram vial and concentrated under nitrogen flow. Toluene (1 mL) was added to aid in removing excess trifluoroacetic acid, and the sample was concentrated again under nitrogen. Lastly methylene chloride (1 mL) was added and the sample was reconcentrated to give 198.3 mg of a golden oil. HPLC (as described earlier, 220 nm) shows a 91% pure major peak. NMR (DMSO): 2.72 (d,  $j=7$  Hz, 2H), 2.79 (s, 6H), 3.87 (d,  $j=7$  Hz, 2H), 5.11-5.20 (m, 1H), 7.30-7.58 (m, 5H), 7.70-7.80 (m, 4H), 8.55 (d,  $j=8$  Hz, 1H), 8.76 (t,  $j=3$  Hz, 1H), 9.39 (s, 1H). MS(ES): product ion observed at  $m/z$  480.

- 567 -

Example 295

Preparation of (±) 3,5-dichloro-β-[[2-[[[3-  
[[ (ethylamino) (methylimino) methyl] amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
benzenepropanoic acid,  
trifluoroacetate salt




The resin product from Step I was treated with 2.5 mL of 95% trifluoroacetic acid/water for 1 hour at 25°C. The filtrate was collected. The resin was washed with 2 x 1 mL of 50% trifluoroacetic acid/methylene chloride and the filtrate collected. The resin was washed once more with 1 mL of methylene chloride. All filtrates were combined into a tared 2 dram vial and concentrated under nitrogen flow. Toluene (1 mL) was added to aid in removing excess trifluoroacetic acid, and the sample was concentrated again under nitrogen. Lastly methylene chloride (1 mL) was added and the sample was reconcentrated to give 261.2 mg of a golden oil. HPLC (as described earlier, 220 nM) shows a 94% pure major peak. NMR (DMSO): 1.11 (t, j=7Hz, 3H), 2.72 (d, J=7 hZ, 2H), 2.79 (s, 3H), 3.25-3.60 (m, 2H), 3.87 (d, j=7 Hz, 2H), 5.02-5.20 (m, 1H), 7.30-7.58 (m, 5H), 7.70-7.85 (m, 4H), 8.55 (d, j=8 Hz, 1H), 8.76 (t, j=3Hz, 1H), 9.40 (s, 1H). MS(ES): product ion observed at m/z 494.



- 568 -


Example 296

Preparation of ( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(1-methylethyl)amino](methylimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid, trifluoroacetate salt



The resin product from Step J was treated with 2.5 mL of 95% trifluoroacetic acid/water for 1 hour at 25°C. The filtrate was collected. The resin was washed with 2 x 1 mL of 50% trifluoroacetic acid/methylene chloride and the filtrate collected. The resin was washed once more with 1 mL of methylene chloride. All filtrates were combined into a tared 2 dram vial and concentrated under nitrogen flow. Toluene (1 mL) was added to aid in removing excess trifluoroacetic acid, and the sample was concentrated again under nitrogen. Lastly methylene chloride (1 mL) was added and the sample was reconcentrated to give 330.3 mg of a golden oil. HPLC (as described earlier, 220 nM) shows an 89% pure major peak. NMR (DMSO): 1.15 (d,  $j=7$ Hz, 6H), 2.72 (d,  $j=7$ Hz, 2H), 2.79 (d,  $j=7$  Hz, 3H), 3.79-3.92 (m, 3H), 5.05-5.20 (m, 1H), 7.30-7.50 (m, 5H), 7.60-7.78 (m, 4H), 8.55 (d,  $j=8$  Hz, 1H), 8.76 (t,  $j=3$ Hz, 1H), 9.40 (s, 1H). MS(ES): product ion observed at  $m/z$  508.

- 569 -

Examples 297-299Step A

10 To a 50 mL round bottom flask equipped with magnetic stirrer was added 3-amino-3-(4-fluoro-phenyl)-propionic acid, (0.300 g, 1.64 mmol). The propionic acid was dissolved in 1 mL of acetone and 6 mL of water and sodium carbonate was added (0.53 g, 4.92 mmol). The pH=10. The

15 Fmoc succinimidyl carbonate (Sigma Chemical Co., 0.553 g, 1.64 mmol) was dissolved in 6 mL of acetone and added slowly to the basic aqueous solution via addition funnel over 20 minutes. The reaction was stirred for 16 hours at room temperature. HPLC analysis (Waters, C18, reverse

20 phase, 25 cm column, 50-90% acetonitrile in water over 30 minutes) indicated that the starting material was consumed. The acetone was removed from the reaction mixture *in vacuo*. The basic aqueous phase was acidified to pH=3 using 3.0 N hydrochloric acid. In a 50 mL

25 separatory funnel the acid layer was washed with 15 mL of ethyl acetate, the water layer was removed and the organic layer was washed (2 x 30 mL water, 2 x 30 mL saturated sodium chloride). The organic layer was dried (magnesium sulfate), filtered and concentrated *in vacuo*. Petroleum


30 ether was added (10 mL) and a white flocculent solid precipitated. After 24 hours of air drying, isolated 0.582 g as a first crop (87.5% yield). The mother liquor was saved for future use. NMR (DMSO): 2.55-2.75 (m, 2H), 4.10-4.30 (m, 3H), 4.85-4.95 (m, 1H), 7.12 (t,  $j=8$  Hz,

- 570 -

2H), 7.24-7.42 (m, 5H), 7.64 (d,  $j=8$  Hz, 2H), 7.82-7.94 (m, 3H). MS (FAB): product ion  $M+Li$  observed at  $m/z$  412.

### Step B

5



10


Wang resin (0.60 g, 0.36 mmol) was placed in a 100 mL round bottom flask. The resin was swelled with 8 mL of methylene chloride for 15 minutes then drained. The FMOC protected amino acid of Step A (0.365 g, 0.9 mmol) was activated in a separate 25 mL round bottom flask by dissolving in methylene chloride/dimethylformamide (4:1, 19 mL) and adding diisopropylcarbodiimide (DIC, 0.141 mL, 0.90 mmol) via syringe, followed by the addition of dimethylaminopyridine (DMAP, 22 mg, 0.18 mmol). The solution was stirred at 25°C for 15 minutes, then added to the preswelled Wang resin. The slurry was stirred for 2 hours at 25°C. The reaction was drained and washed with methanol (3 x 10 mL), methylene chloride (3 x 10 mL) and diethyl ether (3 x 10 mL). To ensure complete reaction, the coupling sequence was repeated. After drying *in vacuo* the resin was swelled with 8 mL of methylene chloride, drained and 8 mL of 20% piperidine/dimethylformamide was added and the slurry was stirred for 30 minutes. The resin was drained and washed as described previously. The resin was dried *in vacuo* for 1 hour. Elemental analysis calculated for resin bound material:

25

30


Calculated: C, 88.23; H, 7.36; N, 0.76; F, 1.03.  
Found: C, 87.13; H, 7.31; N, 0.79; F, 1.06.

- 571 -

Step C

The resin product from Step B was swelled with 8 mL of methylene chloride, then drained. An activated solution of FMOC-Glycine (0.267 g, 0.90 mmol, DIC, 0.140 mL, 0.90 mmol, DMAP, 22 mg, 0.18 mmol. In 10 mL methylene chloride/dimethylformamide, 4:1) was added to the preswelled resin via syringe and stirred at 25°C for 2 hours. The resin was drained and washed (methylene chloride, methanol and diethyl ether, each 3 x 10 mL). The resin was preswelled with 20 mL of methylene chloride for 15 minutes, drained and the coupling reaction was repeated to ensure complete reaction. The Kaiser test (Kaiser, E., Color Test for Detection of Free Terminal Amino Groups in the Solid-Phase Synthesis of Peptides. Anal. Biochem. 1970, 34, 595-598) indicated the coupling was complete. The resin was then suspended in 8 mL of 20% piperidine/dimethylformamide for 30 minutes, drained and washed with dimethylformamide, methanol, methylene chloride, and diethyl ether (3 x 10 ml, each). The resin was dried *in vacuo* for 1 hour.


- 572 -

Step D

5

In a 500 mL round bottom flask equipped with magnetic stirrer, 3-amino-benzoic acid (Aldrich, 10.0 g, 50.8 mmol) was dissolved in 50 mL of dioxane and 133 mL of 10% sodium carbonate. The stirred solution was cooled to 0°C (ice/water) and a solution of fluorenylmethyl chloroformate (13.78 g, 53.3 mmol, in 50 mL dioxane) was added dropwise over 15 minutes. The reaction warmed to 25°C overnight. HPLC analysis (as described earlier) indicated that the starting material was consumed. 500 mL of water was added to the reaction mixture and a white precipitate formed immediately. The solid was collected, washed with 10% citric acid and dried under vacuum. Isolated 15.23 g, (83.4% yield) of a white flocculent solid. NMR (DMSO): 4.18-4.25 (m, 3H), 7.25-7.41 (m, 6H), 7.62-7.72 (m, 3H), 7.80-7.90 (m, 3H). MS (FAB): product ion M+H observed at m/z 360.

20

Step E

25


30

The resin product from Step C was then preswelled using 10 mL of methylene chloride, drained and the activated product of Step D (0.343 g, 0.90 mmol, DIC,

- 573 -

0.141 mL, 0.90 mmol, DMAP, 22 mg, 0.18 mmol, in 5 mL methylene chloride/dimethylformamide 4:1) was added to the preswelled resin. The reaction was stirred for 16 hours at 25°C. The resin was drained and washed as previously described. The Kaiser test indicated that the reaction was not complete. The coupling reaction was repeated, the resin was drained and washed. A repeat Kaiser test indicated that the coupling reaction was complete. The resin was dried in vacuo for 1 hour.

Step F



The resin product from Step E was placed in a 100 mL round bottom flask, was preswelled with 10 mL of dimethylformamide, drained, then treated with 20 mL of 20% piperidine/dimethylformamide for 10 minutes at 25°C. The resin was drained and the procedure was repeated. The resin was filtered and washed with dimethylformamide, methanol, methylene chloride and diethyl ether (3 x 10 mL, each). The Kaiser test indicated that the deprotection step was complete. The resin was placed into a glass 2 dram vial with dimethylformamide (8.0 mL), followed by methyl isothiocyanate (0.526 g, 7.2 mmol). The vial was tightly capped and heated to 80°C for 4 hours. The resin was filtered and washed with dimethylformamide, methanol, methylene chloride, and diethyl ether (3 x 10 mL, each).

- 574 -

The resin was dried in vacuo. Elemental analysis calculated for resin bound material:

Calc'd: C, 83.56; H, 6.46; N, 2.19; F, 1.03; S, 1.35.

Found: C, 82.32; H, 6.67; N, 2.53; F, 1.02; S, 1.44.

5

Step G

The resin product from Step F (100 mg, 0.06 mmol) was transferred to a 2 dram glass vial. The resin was swelled with methylene chloride (3 x 1 mL) and drained. In a separate vial 2-chloro-1-methylpyridiniumiodide (Aldrich, 18.4 mg, 0.072 mmol) was dissolved in 3 mL of dimethylformamide/methylene chloride 4:1 and added to the preswelled resin, followed by triethylamine (20.1 uL, 0.144 mmol). The reaction slurry was stirred for 16 hours at 25°C. The resin was drained, and washed with dimethylformamide, methanol, methylene chloride, and diethyl ether (3 x 4 mL, each). The resin was dried in vacuo for 3 hours. The resin was treated with 95% trifluoroacetic acid (1.5 mL) for 1 hour. The resin was filtered and washed with 50% trifluoroacetic acid/methylene chloride (2 x 1.0 mL) followed by methylene chloride (1 x 1.0 mL). The filtrates were combined and dried in vacuo in tared 2 dram glass vials.

10


15

20

- 575 -

Example 297

Preparation of ( $\pm$ )  $\beta$ -[[2-[[[3-[[[(ethylamino)-(methylimino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]-4-fluorobenzene-  
propanoic acid, trifluoroacetate salt




Isolated 28.1 mg of a golden oil. NMR (DMSO): 1.13 (t,  $j=7$  Hz, 3H), 2.65-2.75 (m, 2H), 2.76-2.85 (m, 3H), 3.25 (t,  $j=3$  Hz, 2H), 3.80-3.95 (m, 2H), 5.10-5.21 (m, 1H), 7.13 (t,  $j=8$  Hz, 2H), 7.30-7.40 (m, 3H), 7.52 (t,  $j=8$  Hz, 1H), 7.65-7.85 (m, 3H), 8.49 (d,  $j=8$  Hz, 1H), 8.71 (t,  $j=8$  Hz, 1H) 9.40 (s, 1H). HPLC (as described earlier, 220 nm) 90.15% pure. MS (ES): product ion observed at  $m/z$  444.



- 576 -

Example 298

Preparation of ( $\pm$ ) 4-fluoro- $\beta$ -[[2-[[[3-[[[(1-methylethyl)amino](methylimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid, trifluoroacetate salt




Isolated 44.9 mg of a golden oil. NMR (DMSO): 1.16 (d,  $j=7$  Hz, 6H), 2.61-2.70 (m, 2H), 2.73-2.80 (m, 3H), 3.75-3.90 (m, 3H), 5.10-5.21 (m, 1H), 7.11 (t,  $j=8$  Hz, 2H), 7.25-7.37 (m, 3H), 7.49 (t,  $j=8$  Hz, 1H), 7.59-7.82 (m, 3H), 8.49 (d,  $j=8$  Hz, 1H), 8.70 (t,  $j=3$  Hz, 1H) 9.40 (s, 1H). HPLC (as described earlier, 220 nm) 98% pure. MS (ES): product ion observed at  $m/z$  458.

- 577 -


Example 299

Preparation of (±) 4-fluoro-β-[[2-[[[3-[[[(4-pyridinylmethyl)amino](methylimino)methyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid, trifluoroacetate salt










Isolated 31.6 mg of a golden oil. NMR (DMSO): 2.60-2.72 (m, 2H), 2.81-2.89 (d,  $j=7$  Hz, 3H), 3.80-3.95 (m, 2H), 4.61-4.80 (bs, 2H), 5.10-5.21 (m, 1H), 7.01-7.22 (m, 4H), 7.29-7.44 (m, 3H), 7.50 (t,  $j=8$  Hz, 1H), 7.65-7.85 (m, 3H), 8.40-8.50 (d,  $j=8$  Hz, 1H), 8.70-8.85 (m, 3H), 9.73 (s, 1H). HPLC (as described earlier, 220 nm) 98% pure. MS(ES): product ion observed at  $m/z$  507.








The following compounds are prepared according to analogous solid-phase synthetic methods described in Examples 294-299.










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
300	Cl	H	Cl	-H	
301	Cl	H	Cl	-H	
302	Cl	H	Cl	-H	
303	Cl	H	Cl	-H	











- 579 -

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
304	Cl	H	Cl	-H	
305	Cl	H	Cl	-H	
306	Cl	H	Cl	-H	-CH <sub>2</sub> CH <sub>3</sub>
307	Cl	H	Cl	-H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>
308	Cl	H	Cl	-H	
309	Cl	H	Cl	-H	
310	Cl	H	Cl	-H	
311	Cl	H	Cl	-H	
312	Cl	H	Cl	-H	



Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
313	Cl	H	Cl	-H	
314	Cl	H	Cl	-H	
315	Cl	H	Cl	-H	
316	Cl	H	Cl	-H	
317	Cl	H	Cl	-H	
318	Cl	H	Cl	-H	
319	Cl	H	Cl	-H	

- 581 -

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
320	Cl	H	Cl	-CH <sub>3</sub>	
321	Cl	H	Cl	-CH <sub>3</sub>	 .TFA
322	Cl	H	Cl	-CH <sub>3</sub>	 .TFA
323	Cl	H	Cl	-CH <sub>3</sub>	 .TFA
324	Cl	H	Cl	-CH <sub>3</sub>	
325	Cl	H	Cl	-CH <sub>3</sub>	 .TFA
326	Cl	H	Cl	-CH <sub>3</sub>	-CH <sub>2</sub> (CF <sub>2</sub> ) <sub>2</sub> CF <sub>3</sub>
327	Cl	H	Cl	-CH <sub>3</sub>	

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
328	Cl	H	Cl	-CH <sub>3</sub>	
329	Cl	H	Cl	-CH <sub>3</sub>	
330	Cl	H	Cl	-CH <sub>3</sub>	
331	Cl	H	Cl	-CH <sub>3</sub>	
332	Cl	H	Cl	-CH <sub>3</sub>	
333	Cl	H	Cl	-CH <sub>3</sub>	
334	Cl	H	Cl	-CH <sub>3</sub>	
335	Cl	H	Cl	-CH <sub>3</sub>	
336	Cl	H	Cl	-CH <sub>3</sub>	
337	Cl	H	Cl	-CH <sub>3</sub>	

- 583 -


Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
338	Cl	H	Cl	-CH <sub>3</sub>	
339	Cl	H	Cl	-CH <sub>3</sub>	



- 584 -

Example 361

Preparation of (±) ethyl 3,5-dichloro-β-[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-  
5 amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate, trifluoroacetate salt


Step A

To a suspension of the 1-(3-carboxyphenyl)-2-thiourea  
15 (produced in Example 236, Step A) (10.00 g, 0.051 mol) in  
ethanol (100 mL) was added iodomethane (3.5 mL) and heated  
at 70°C under nitrogen atmosphere for 2.5 hours. The  
reaction mixture was concentrated under reduced pressure,  
the residue was triturated with ether containing 10% EtOAc  
20 (2 x 100 mL) and the supernatant decanted. The resulting  
solid was dried in vacuo for 2 hours, dissolved in DMF (75  
mL) and added dropwise to a solution of 2,2 dimethyl-1,3  
propanediamine (42 g, 0.41 mol) in DMF (20 mL) over a  
period of 1 hour. The resulting mixture was heated at  
25 80°C under nitrogen atmosphere for 16 hours with  
simultaneous trapping of the methylmercaptan in 5% sodium  
hypochlorite solution. DMF was distilled in vacuo, the  
residue was dissolved in water (50 mL) and washed with  
diethyl ether (3 x 25 mL). The aqueous phase was  
30 acidified with 2N HCl to pH 4.0 when a white precipitate  
was obtained. It was filtered, washed with water and  
ether and dried to give the desired product 8.0 g (63%)  
as a white powder. <sup>1</sup>H-NMR and MS were consistent with the  
structure.

- 585 -

Step B

5




To a suspension of the HCl salt of Step A (1.0 g, 0.0035 mol) in DMF (15 mL), was added N-methylmorpholine (0.46 mL) and cooled to -10°C in an ice-salt bath. This reaction mixture was then treated with isobutyl chloroformate (0.45 mL), stirred at -10°C for 30 minutes, and a solution of the amine generated by the addition of N-methylmorpholine (0.46 mL) to a solution of t-butylglycinate hydrochloride (0.6 g) in DMF (5 mL) at 0°C. The resulting reaction mixture was stirred at -10°C for 1 hour and at room temperature for 16 hours under argon atmosphere. DMF was distilled in vacuo, the residue was treated with 5% sodium bicarbonate (25 mL) and EtOAc (25 mL) and stirred at room temperature for 30 minutes. A white precipitate was obtained. The precipitate was filtered, washed with water (2 x 20 mL), and EtOAc (2 x 20 mL), and dried to give the desired compound, 0.58 g (46%). <sup>1</sup>H-NMR and MS were consistent with the structure.

25


Step C

30



- 586 -

The product of Step B (0.6 g, 0.0017 mol) was suspended in dioxane (2.0 mL) and treated with 4N HCl in dioxane (0.9 mL) and stirred overnight at room temperature. The reaction mixture was diluted with diethyl ether, filtered, and the residue washed with diethyl ether (3 x 20 mL). The resulting pale yellow solid was dried in a desiccator over NaOH pellets and used as such in the following step, without purification.

10 Step D

To a suspension of HCl salt as prepared in Step C in DMF (10 mL), was added N-methylmorpholine (0.21 mL) and cooled to -10°C in an ice-salt bath. This reaction mixture was then treated with isobutylchloroformate (0.24 mL), stirred at -10°C for 30 minutes, and a solution of the amine generated by the addition of N-methylmorpholine (0.46 mL) to a solution of ethyl DL-3-amino-3-(3,5-dichlorophenyl)-propionate (produced as in Example 1, Steps A and B substituting 3,5-dichlorobenzaldehyde for 3-pyridine carboxaldehyde) (0.6 g, 0.002 mol) in DMF (5 mL) at 0°C. The resulting reaction mixture was stirred at -10°C for 1 hour and at room temperature for 16 hours under argon atmosphere. DMF was distilled in vacuo, the residue was triturated with ether (2 x 25 mL) and the supernatant decanted. The insoluble residue was purified by reverse phase HPLC using a 30 minute gradient of 5-70% CH<sub>3</sub>CN in water at a flow rate of 70 mL/minute. The appropriate

- 587 -


fractions were combined and freeze dried to afford the desired TFA salt, as a pale yellow powder. <sup>1</sup>H-NMR and MS were consistent with the structure.

- 588 -

Example 362

Preparation of (±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-  
5 amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid, trifluoroacetate salt

10




15

The above compound was prepared by coupling the  
product of Step C in Example 361 with the product of  
Example 440, Step A, as described in Example 361. The  
20 desired product was isolated by reverse-phase HPLC using a  
30 minute gradient of 5-70% CH<sub>3</sub>CN in water at a flow rate  
of 70 mL/minute. The appropriate fractions were combined  
and freeze dried to afford the desired TFA salt. <sup>1</sup>H-NMR  
and MS were consistent with the structure.

- 589 -

Example 363

Preparation of  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
5 fluorobenzenepropanoic acid,  
trifluoroacetate salt

Step A

15 A mixture of 4-fluorophenyl bromide (10.0 g, 0.057 mol), tert-butylacrylate (9.52 g, 0.074 mol), palladium acetate (0.13 g, 0.00057 mol), tri-para-tolylphosphine (0.87 g, 0.0029 mol) and triethylamine (5.78 g, 0.057 mol) in 30 mL of DMF was heated at 120°C for 16 hours. The  
20 mixture was cooled and treated with 500 mL of water. The aqueous phase was extracted with ethyl acetate (3 x 200 mL) and the organic layer was washed with brine, dried over magnesium sulfate and filtered. The filtrate was concentrated and the residue was purified by  
25 chromatography on silica gel (ethyl acetate/hexane, 1:9) to give 10.13 g of product as a yellow oil (80%). The NMR was consistent with the proposed structure.

Analysis Calc'd. for  $C_{13}H_{15}FO_2$ : C, 70.25; H, 6.80.

Found: C, 69.77; H, 7.08.

30

- 590 -

Step B

The product from Step A (8.7 g, 0.039 mol) was treated with tert-butanol saturated with ammonia and 3 mL of acetic acid at 110°C and 900 psi in a Parr shaker for 5 48 hours. The mixture was filtered and concentrated. The residue was dissolved with 200 mL of cold 1N HCl and extracted with ethyl acetate. The aqueous phase was then basified with potassium carbonate and extracted with methylene chloride (2 x 200 mL). The combined extracts 10 were washed with brine, dried over magnesium sulfate and filtered. The filtrate was concentrated to give 4.23 g of a yellow oil (41%). The structural assignment was supported by the NMR spectrum.

15 Step C

To a solution of the compound of Example M (1.0 g, 0.0037 mol) in 10 mL of DMF was added N-methylpiperidine (0.42 g, 0.0037 mol) rapidly. The mixture was stirred at room temperature for 20 minutes, then treated with 20 isobutyl chloroformate at 0°C. After 15 minutes, a solution of the product from Step B in 3 mL of DMF was added. The reaction mixture was allowed to warm to room temperature and stirred overnight. Dimethylformamide was removed in vacuo and the residue was purified by reverse 25 phase HPLC [acetonitrile/water (containing 0.5% of TFA)] to give 0.97 g of a pale yellow solid (44%):

Analysis Calc'd. for  $C_{23}H_{27}N_3O_4F \cdot 1.0 H_2O \cdot 1.0 TFA$ :

C, 50.93; H, 5.30; N, 11.88.

Found: C, 50.61; H, 4.92; N, 11.74.

30

Step D

To a suspension of the product from Step C in 10 mL of methylene chloride at 0°C was added 6 mL of TFA. The mixture was stirred at room temperature for 4 hours..

- 591 -

Solvent was removed and the residue was purified by reverse phase HPLC [acetonitrile/water (containing 0.5% of TFA)] to give 0.75 g of the title compound as a white solid (75%):

5 Analysis Calc'd. for  $C_{19}H_{20}N_3O_4F \cdot 1.5$  TFA:

C, 46.16; H, 3.79; N, 12.23.


Found: C, 45.86; H, 3.68; N, 12.23.



- 592 -

Example 364

Preparation of ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1H-imidazole-2-propanoic acid,  
tris(trifluoroacetate) salt

Step A

A solution of 2-imidazolecarboxaldehyde (6.0 g, 0.063 mol) and (tert-butylcarbonylmethylene)triphenylphosphorane (29.4 g, 0.078 mol) in 150 mL of tetrahydrofuran was heated at 55°C overnight. The clear solution was cooled and concentrated *in vacuo*. The residue was purified by chromatography on silica gel (ethyl acetate/hexane, 8:2) to give 9.7 g of product (1:1 E/Z mixture) as a white solid (79%): Analysis Calc'd. for  $C_{10}H_{14}N_2O_2$ :

C, 61.84; H, 7.27; N, 14.42.

Found: C, 61.52; H, 7.39; N, 14.21.

Step B

To a suspension of prewashed sodium hydride (0.62 g, 0.026 mol) in 40 mL of dry dimethylformamide was added the product from Step A slowly. After 30 minutes, 2-(trimethylsilyl)ethoxymethyl chloride was added and the reaction mixture was stirred at room temperature for 2 hours. Water was added and the aqueous phase was extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and filtered. The filtrate was concentrated and the residue

- 593 -

purified by chromatography on silica gel (ethyl acetate/hexane, 1:1) to give 3.54 g of E isomer as a colorless oil and 2.66 g of Z isomer as a white solid (73%). Analysis Calc'd. for  $C_{16}H_{27}N_2O_2Si$ :

5 C, 59.22; H, 8.70; N, 8.63.

**Found:** C, 58.94; H, 9.12; N, 8.53.

**Step C**

To a solution of N-benzyl(trimethylsilyl)amine (2.16 g, 0.012 mol) in 30 mL of dry tetrahydrofuran at -78°C was added n-butyllithium (0.012 mol) slowly. After 30 minutes, a solution of the product of Step B (2.6 g, 0.008 mol) in 15 mL of tetrahydrofuran was added and the reaction mixture was stirred at this temperature for 2.5 hours. The reaction was then quenched with a solution of acetic acid in tetrahydrofuran, followed by addition of saturated sodium bicarbonate to pH 9. The aqueous phase was extracted with ethyl acetate and the organic layer was washed with brine, dried over magnesium sulfate and filtered. The filtrate was concentrated and purified by chromatography on silica gel (ethyl acetate/hexane, 6:4) to give 1.96 g of product as a clear oil (60%).

**Analysis Calc'd. for  $C_{23}H_{27}N_3O_3Si$ :**

C, 64.00; H, 8.64; N, 9.73.


25 Found: C, 63.72; H, 8.85; N, 9.73.

**Step D**

To a solution of the product from Step C (5.4 g, 0.0125 mol) and ammonium formamide (7.89 g, 0.125 mol) in 150 mL of methanol was added Pd/C (170 mg). The mixture was stirred at reflux for 3 hours. The catalyst was filtered through celite and the filtrate was concentrated. The residue was dissolved in 400 mL of water, saturated with potassium carbonate, extracted with ethyl acetate.

- 594 -

The organic layer was washed with brine, dried over magnesium sulfate and filtered. The filtrate was concentrated to give 3.9 g of product as a colorless oil (91%). The NMR spectrum indicated that the compound was of sufficient purity for the next step.

Step E

The above compound was synthesized under the same conditions as described in Step C of Example 363. The crude product was purified by reverse phase HPLC [acetonitrile/water (containing 0.5% of TFA)] to give 1.5 g of product as a yellow solid (60%):

20 Analysis Calc'd. for  $C_{26}H_{41}N_7O_5Si \cdot 2.5$  TFA:

C, 44.07; H, 5.19; N, 11.61.

Found: C, 44.24; H, 5.14; N, 11.91.

Step F

25 The title compound was obtained from the product of Step E following the procedure described in Step D of Example 363. The crude product was purified by reverse phase HPLC [acetonitrile/water (containing 0.5% of TFA)] to give 0.35 g of the title compound as a yellow solid (24%):

30 Analysis Calc'd. for  $C_{16}H_{19}N_7O_4 \cdot 3.0$  TFA:

C, 36.93; H, 3.10; N, 13.70.


Found: C, 37.76; H, 2.95; N, 14.22.

- 595 -

Example 365

Preparation of (±) β-[[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
2,3,5,6-tetrafluorobenzenepropanoic acid,  
trifluoroacetate salt

10



15

The above compound was made by following the reaction  
sequence described in Example 364 Step A and Step C to  
Step F. The structure was confirmed by the NMR spectrum.  
Analysis Calc'd. for  $C_{19}H_{17}N_5O_4F_4 \cdot 1.5$  TFA:

C, 42.18; H, 2.98; N, 11.18.


Found: C, 42.24; H, 3.07; N, 11.12.

20

- 596 -

Example 366

Preparation of  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
5 -5-bromothiophene-2-propanoic acid,  
trifluoroacetate salt, monohydrate



15

A solution of the product of t-butyl ester of the  
above compound (prepared according to analogous  
methodology as described herein) (1.0 g, 1.91 mmol) and  
20 trifluoroacetic acid (14.8 g, 10.0 ml, 13.0 mmol) in  
dichloromethane (25 ml) was stirred at 0°C for 30 minutes.  
The reaction mixture was allowed to warm to room  
temperature and stirred for 6 hours. The solvent was  
removed under reduced pressure. The crude product was  
25 purified by HPLC (acetonitrile, water, trifluoroacetic  
acid) to give pure title compound (0.43 g, 38%) as a white  
solid.


Analysis Calc'd. for  $C_{17}H_{18}N_4O_4SBr \cdot CF_3COOH \cdot H_2O$ :

30 Found: C, 38.01; H, 3.53; N, 11.67; S, 5.34  
C, 38.07; H, 3.23; N, 11.48; S, 4.99

- 597 -

Example 367

Preparation of ( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-  
2-yl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid,  
trifluoroacetate salt




The ethyl ester prepared in Example 361, Step D (0.22 g) was hydrolyzed to the acid using 1M LiOH, (1.8 mL) in acetonitrile (0.2 mL), followed by acidification and purification by reverse-phase HPLC to give 0.18 g of the acid as pale yellow powder.  $^1\text{H}$  NMR and MS were consistent with the structure.

- 598 -

Example 368

Preparation of ( $\pm$ ) 3-bromo-5-dichloro-2-hydroxy-  
 $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-  
5 dimethylpyrimidin-2-yl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]benzene-  
propanoic acid, trifluoroacetate salt




The above compound was prepared by coupling the acid  
prepared in Example 361, Step C, (0.6 g) with the product  
20 of Example 233, Step B (0.5 g) according to the procedure  
described in Example 361. The desired product was  
isolated by reverse-phase HPLC to give 0.38 g of the above  
compound as a pale yellow powder.  $^1\text{H}$  NMR and MS were  
consistent with the structure.

25

- 599 -

Example 370

Synthesis of  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-2-  
mercaptobenzenepropanoic acid, lithium salt

Step A

Synthesis of S-Phenyl Thiocinnamate: A solution of cinnamoyl chloride (14.6 g, 87.68 mmol) in dichloromethane (100 mL) was added to a solution of thiophenol (9.55 g, 86.68 mmol) and pyridine (7 mL) in dichloromethane (150 mL) in an ice-water bath. After 18 hours at room temperature, the reaction mixture was washed with dilute hydrochloric acid (100 mL, 1N), brine (100 mL), dried ( $\text{MgSO}_4$ ) and was concentrated to afford 19.0 g (91%) of the desired thioester as a crystalline solid.

Step B

Synthesis of Thiocoumarin: A mixture of S-phenyl thiocinnamate (14.0 g, 58.25 mmol) and aluminum chloride (39 g) was stirred and heated at 85°C for 3 hours. The hot reaction mixture was poured carefully over ice, then was extracted with ethyl acetate (3 x 300 mL), washed with brine (200 mL), dried ( $\text{MgSO}_4$ ) and was concentrated. The residue was recrystallized from hexane-ethyl acetate to afford 5.2 g (52%) of the desired product as pale yellow crystals.



- 600 -

Step C

## Synthesis of 4-Amino-3,4-Dihydrothiocoumarin

Hydrochloride Salt: Lithium hexamethyldisilazane (10.22 mL, 1N, 10.22 mmol) was added slowly to a solution of thiocoumarin (1.41 g, 8.52 mmol) in tetrahydrofuran (20 mL) at -78°C. After 45 minutes, the reaction mixture was warmed up to 0°C, then was quenched with glacial acetic acid (0.511 g). After 10 minutes, the reaction mixture was partitioned between ethyl acetate (100 mL) and sodium bicarbonate (100 mL). The organic layer was dried (MgSO<sub>4</sub>) and was concentrated. The residue obtained was dissolved in ether (100 mL) and dioxane/HCl (20 mL, 4N) was added. The precipitate formed was filtered and the solid was dried in vacuo to afford (0.50 g, 27%) of the desired product as a yellow powder.

Step D


A solution of m-guanidinohippuric acid (0.506 g, 1.855 mmol) in dimethylformamide (5 mL) and N-methylmorpholine (0.187 g, 1.855 mmol) was cooled to 0°C and was stirred for 15 minutes. Isobutylchloroformate (0.253 g, 1.855 mmol) was added in three portions. After 10 minutes, 4-amino-3,4-dihydrothiocoumarin hydrochloride (0.404 g, 1.855 mmol) was added in one portion followed by N-methylmorpholine (0.187 g, 1.855 mmol). The reaction mixture was stirred for 18 hours at room temperature. The reaction mixture was concentrated and the residue was dissolved in tetrahydrofuran/water (1:1, 5 mL) and was chromatographed (reverse phase, 95:5 water: acetonitrile over 60 minutes to 30:70 water: acetonitrile containing 0.1% TFA). The eluents were lyophilized to afford 0.300 g of the title compound as a pale yellow powder.

Proton NMR and MS were consistent with the desired product.

- 601 -

Example 371

Preparation of ( $\pm$ )  $\beta$ -[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-5-  
chloro-2-mercaptobenzenepropanoic acid,  
dilithium salt

Step A

Synthesis of S-(4-Chlorophenyl) Thiocinnamate: A solution of cinnamoyl chloride (26.0 g, 156.3 mmol) in dichloromethane (100 mL) was added to a solution of thiophenol (22.6 g, 156.3 mmol) and pyridine (12.6 mL) in dichloromethane (200 mL) in an ice-water bath. After 18 hours at room temperature, the reaction mixture was washed with dilute hydrochloric acid (100 mL, 1N), brine (100 mL), dried ( $\text{MgSO}_4$ ) and was concentrated to afford 41.0 g (96%) of the desired thioester as a crystalline solid.

Step B

Synthesis of 6-Chlorothiocoumarin: A powdered mixture of S-(4-chlorophenyl) thiocinnamate (19.4 g) and aluminum chloride (52 g) was stirred and heated at 125°C for 3 hours. The hot reaction mixture was poured carefully over ice/water, then was extracted with ethyl acetate (3x300 mL), washed with brine (200 mL), dried ( $\text{MgSO}_4$ ) and was concentrated. The residue was triturated

- 602 -

with hexane/ethyl acetate to afford 2.0 g (14%) of the desired product as pale yellow crystals.

### Step C

5        Synthesis of 4-amino-6-chloro-3,4-dihydrothi coumarin hydrochloride salt: Lithium hexamethyldisilazane (6.4 mL, 1N, 6.4 mmol) was added slowly to a solution of  
10       6-chlorothi coumarin (1.05 g, 5.345 mmol) in tetrahydrofuran (20 mL) at -78°C. After 45 minutes, the  
15       reaction mixture was warmed up to 0°C, then was quenched with glacial acetic acid (0.321 g). After 10 minutes, the  
     reaction mixture was partitioned between ethyl acetate (100 mL) and sodium bicarbonate (100 mL). The organic  
     layer was dried (MgSO<sub>4</sub>) and was concentrated. The residue  
     obtained was dissolved in ether (100 mL) and dioxane/HCl (20 mL, 4N) was added. The precipitate formed was  
     filtered and the solid was dried in vacuo to afford  
     (0.80 g, 60%) of the desired product as a yellow powder.

### 20      Step D

     A solution of m-guanidinohippuric acid (0.548 g, 2.0 mmol) in dimethylformamide (5 mL) and N-methylmorpholine (0.220 mL, 2.0 mmol) was cooled to 0°C and was stirred for  
25       15 min. Isobutylchloroformate (0.260 mL, 2.0 mmol) was  
     added in three portions. After 10 minutes, 4-amino-6-chloro-3,4-dihydrothi coumarin hydrochloride (0.50 g, 2.0 mmol) was added in one portion followed by N-methylmorpholine (0.220 mL, 2.0 mmol). The reaction  
30       mixture was stirred for 18 hours at room temperature. The  
     reaction mixture was concentrated and the residue was  
     dissolved in tetrahydrofuran/water (1:1, 5 mL) and was  
     chromatographed (reverse phase, 95:5 water: acetonitrile over 60 minutes to 30:70 water: acetonitrile containing 0.1% TFA). The eluents were basified with an aqueous

- 603 -

solution of lithium hydroxide and then was lyophilized to afford 0.300 g of the title compound as a pale yellow powder.

MS and NMR were consistent with the proposed  
5 structure.


- 604 -

Example 372

The following compounds are prepared according to the methodology described in Examples 370-371.

5

10



15

X=SH; R<sub>1</sub>, R<sub>2</sub>=Cl; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>, R<sub>2</sub>=F; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>, R<sub>2</sub>=Me; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>, R<sub>2</sub>=CF<sub>3</sub>; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>, R<sub>2</sub>=Br; R<sub>3</sub>, R<sub>4</sub>=H


20

X=SH; R<sub>1</sub>=H, R<sub>2</sub>=F; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>=H, R<sub>2</sub>=Br; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>=H, R<sub>2</sub>=CF<sub>3</sub>; R<sub>3</sub>, R<sub>4</sub>=HX=SH; R<sub>1</sub>=H, R<sub>2</sub>=CH<sub>3</sub>; R<sub>3</sub>, R<sub>4</sub>=H

25 and the above compounds wherein R<sub>3</sub> and R<sub>4</sub> together are (CH<sub>2</sub>)<sub>3</sub> or (CH<sub>2</sub>)<sub>2</sub>.


- 605 -

### EXAMPLE 374




10 The above compound is prepared by reacting the  
compound prepared in Example 233, Step B with 3-guanidino-  
5-trifluoromethylhippuric acid (prepared according to the  
procedure of Example 38) using substantially the  
proportions and procedure of Example N, Step 3 and  
15 substituting 3-guanidino-5-trifluoromethylhippuric acid  
hydrochloride for GIHA HCl. The desired product is  
isolated by C-18 RPHPLC.

### EXAMPLE 375




The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-6,8-dichloro-hydrocoumarin hydrochloride prepared in Example 237 for the compound of Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

- 606 -

EXAMPLE 376

10 The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-6-chloro-hydrocoumarin hydrochloride prepared in Example 231 for the compound of Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

15

EXAMPLE 377


25

The above compound is prepared using the procedure of Example 374 and substituting the compound prepared in Example 227 for the compound of Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

- 607 -

EXAMPLE 378

5




10 The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-6-nitro-hydrocoumarin hydrochloride prepared in Example 226 for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

15

EXAMPLE 379


20




25 The above compound is prepared from the product of Example 378 using the conditions of Example 234. The desired product is isolated by C-18 RPHPLC.



- 608 -


EXAMPLE 380

The above compound is prepared using the procedure of Example 374 and substituting the compound prepared in Example 235, Steps A-C and two equivalents of NMM in the coupling step for the compound prepared in Example 233, Step B and one equivalent of NMM. The desired product is isolated by C-18 RPHPLC.


EXAMPLE 381

The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-6-methylhydrocoumarin hydrochloride (prepared in Example 88) for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

- 609 -


EXAMPLE 382

The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-hydrocoumarin hydrochloride (prepared in Example 87) for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

EXAMPLE 383

The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-7-methoxy-hydrocoumarin hydrochloride (prepared in Example 222) for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

- 610 -


EXAMPLE 384

10


The above compound is prepared using the procedure of Example 374 and substituting (RS)-4-amino-8-methoxy-hydropsoralen hydrochloride (prepared in Example 223) for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

15

- 611 -

EXAMPLE 385Step A

10 Preparation of



The above compound is prepared from 7,8-methylenedioxy-coumarin (which may be prepared from 7,8-dihydroxy-chromen-2-one according to P. Castillo, J.C. Rodriguez-Ubis, and F. Rodriguez, *Synthesis*, 10, 839-840 (1986)) using the procedure of Example 233, Steps A and B.

Step B


25 The above Example compound is prepared using the procedure of Example 374, substituting the hydrochloride salt of the product of Step A for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

30

- 612 -


EXAMPLE 386

5

Step A

10 Preparation of

15




20 The above compound is prepared from 6,7-methylenedioxy-coumarin [which may be prepared from 6,7-dihydroxy-chromen-2-one according to Spaeth, et al., Chem. Ber., 70, 702 (1937)] using the procedure of Example 233, Steps A and B.

25 Step B


The above Example compound is prepared using the procedure of Example 374 and substituting the hydrochloride salt of the product of Step A for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

30

- 613 -

EXAMPLE 387Step A

10 Preparation of




20 The above compound is prepared from 5,6-methylenedioxy-coumarin [prepared from 5,6-dihydroxy-chromen-2-one according to P. Castillo, J.C. Rodriguez-Ubis, and F. Rodriguez, *Synthesis* 10, 839-840 (1986)] using the procedure of Example 233, Steps A and B.


25 Step B

The above Example compound is prepared using the procedure of Example 374, substituting the hydrochloride salt of the hydrochloride salt of the product of Step A for the compound prepared in Example 233, Step B. The  
30 desired product is isolated by C-18 RPHPLC.

- 614 -

EXAMPLE 388Step A

Preparation of




25 The above compound may be prepared by reacting  
esculin (Aldrich, rendered substantially free from water  
of hydration by storage of  $P_2O_5$  in a vacuum dessicator)  
according substantially to the procedure of S. Kato, et  
al., *Bull. Chem. Soc. Jap.*, 54, 6, 1981, 1895-1896, for  
the conversion of phenyl- $\alpha$ -D-glucoparanoside to phenyl  
30 2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucoparanoside and  
substituting the appropriate molar quantities of reagents  
to effect complete conversion of esculin to the above  
compound. The desired product may be isolated by standard  
silica gel chromatography or by preparative C-18 RPHPLC.

- 615 -

Step B

5

10




The above compound is prepared using the procedure of  
Example 233, Step B and substituting the product of Step A  
15 for the product of Example 233, Step A.

Step C

20


25



The above compound is prepared using the procedure of  
30 Example 374, substituting the hydrochloride salt of the  
product of Step B for the compound prepared in Example  
233, Step B. The desired product is isolated by C-18  
RPHPLC.




- 616 -


Step D

The above compound is prepared by taking the product of Step C, dissolving in a suitable solvent (e.g. aqueous ethanol), transferring to a Fischer-Porter pressure bottle equipped with an inlet and outlet valve, pressure gauge and pressure relief valve and removing the benzyl groups by standard catalytic hydrogenolysis procedure: 5% Pd on carbon catalyst and hydrogen atmosphere until the debenzylation reaction is substantially complete. The desired product is isolated by C-18 RPHPLC.

- 617 -

EXAMPLE 389Step A

10 Preparation of




The above compound is prepared using substantially the procedure of Example 235, Steps A-C.


Step B

The above Example compound is prepared using substantially the procedure of Example 235, Steps D and E and is isolated using preparative C-18 RPHPLC.

- 618 -

EXAMPLE 390Step A


10 Preparation of 4-chloro-2-iodophenol



The above compound is prepared according to the  
procedure of K.J. Edgar and S.N. Felling, *J. Org. Chem.*,  
20 55, 16, 1990, 5287-5291.

Step B

25 Preparation of 5-chloro-3-iodosalicylaldehyde




4-chloro-2-iodophenol prepared in Step A is converted  
to the salicylaldehyde using the procedure of  
G. Casiraghi, et al., *J.C.S. Perkin I*, 1978, 318-321.

- 619 -

**Step C**


### Preparation of 6-chloro-8-iodocoumarin



5-chloro-3-iodosalicylaldehyde is converted into the corresponding coumarin, 6-chloro-8-iodocoumarin, using substantially the procedure of Example 233, Step A and substituting 5-chloro-3-iodo-salicylaldehyde for 3-bromo-15 5-chlorosalicylaldehyde. The desired product may be isolated by standard silica gel chromatography or distillation.

**Step D**

20 Preparation of (R,S)-4-amino-6-chloro-8-iodo-hydrocoumarin



The above compound is prepared using substantially  
30 the procedure of Example 233, Step B and substituting the  
product of Step C for 3-bromo-5-chlorosalicylaldehyde to  
give the product as substantially pure hydrochloride salt.


- 620 -

Step E

The above Example compound is prepared using the procedure of Example 274 and substituting the product of Step D for the compound prepared in Example 233, Step B.


5 The desired product is isolated by C-18 RPHPLC.

- 621 -


EXAMPLE 391

10 The above compound is prepared using substantially the procedure of Example 86, Step D, substituting 3-guanidino-5-trifluoromethylhippuric acid hydrochloride for GIHA HCl. The desired product is isolated by C-18 RPHPLC.

- 622 -

EXAMPLE 392Step A

Preparation of




15

The above compound is prepared using substantially the procedure of Example 235, Step A, substituting BOC-L-aspartic acid-4-tert-butyl ester (Fluka) for 5-bromonicotinic acid.

20

Step B

Preparation of



30

The above compound is prepared according to substantially the procedure of M.R. Angelastro, et al., *J. Med. Chem.*, 1994, 37, 4538-4554, substituting the product of Step A for Reference compound 2 {(S)-[1-

- 623 -

[methoxymethylamino)carbonyl]-2-methylpropyl carbamic acid,  
1,1-dimethylethyl ester} and deprotecting according to  
substantially the procedure employed for obtaining  
reference compound 3 to obtain the above compound as the  
HCl salt.


#### Step C

The above Example compound is prepared using  
substantially the procedure of Example 85, Step A,  
substituting the product of Step B for glycine t-butyl  
ester and substituting 3-guanidino-5-  
trifluoromethylhippuric acid hydrochloride for GIHA HCl.  
The desired product may be obtained by C-18 RPHPLC.

15



- 624 -


EXAMPLE 393Step A

- 10 Preparation of 3-N-t-Boc-amino-4-hydroxy-(3S)-butyric acid benzyl ester

N-t-Boc-L-aspartic acid,  $\beta$ -benzyl ester (10.0 mmole) was dissolved in 10 mL of THF and added dropwise over a period of 30 minutes to a 0°C solution of  $\text{BH}_3$ -THF (20 mL, 20.0 mmole), under argon. After the mixture was stirred for an additional 1-2 hours at 0°C, the reaction was quenched by dropwise addition of 10% acetic acid in methanol and the solvent evaporated. The oil residue was dissolved in ethyl acetate and extracted with 1N HCl, water, and 1M  $\text{NH}_4\text{HCO}_3$ . The ethyl acetate layer was dried ( $\text{Na}_2\text{SO}_4$ ) and volatiles evaporated to give an oil was crystallized from isopropanol/hexane (mp 56-57°C):  $^1\text{H}$  NMR,  $\text{CDCl}_3$ ,  $\delta$ , 1.45 (s, 9H), 2.65 (d, 2H), 3.68 (d, 2H), 5.12 (s, 2H), 5.25 (m, 1H), 7.35 (m, 5H).

Step B

Preparation of



- 625 -

The 3-N-t-Boc-amino-4-hydroxy-butyric acid benzyl ester prepared in Step A was oxidized to the corresponding aldehyde using the following Swern oxidation conditions: oxalyl chloride (6.40 g, 20.72 mmole) was dissolved in dry  $\text{CH}_2\text{Cl}_2$  (25 mL) under argon and cooled to  $-63^\circ\text{C}$  using a dry ice/chloroform bath. Dry DMSO (g, 41.4 mmole) dissolved in  $\text{CH}_2\text{Cl}_2$  (12 mL) was added in a dropwise fashion over 15 minutes. The alcohol (6.40 g, 20.7 mmole), dissolved in methylene chloride (50 mL) was then added over 10 minutes. After stirring the reaction mixture for an additional 10 minutes,  $\text{Et}_3\text{N}$  (11.6 mL, 82.9 mmole, 4.0 equivalents) in  $\text{CH}_2\text{Cl}_2$  (25 mL) was added over 15 minutes. The resulting mixture was stirred for 15 minutes and quenched by addition of water (31 mL). The resulting slurry was poured onto hexanes (250 mL) and the organic layer washed with aqueous  $\text{KHSO}_4$ . The aqueous layer was extracted with diethyl ether and the combined organic extracts were washed with saturated  $\text{NaHCO}_3$ , dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to give 5.8 g of a light yellow oil which was substantially the desired aldehyde. A small portion was purified by flash chromatography (hexane: ethyl acetate, Merck 60 silica gel):  $^1\text{H}$  NMR (300 MHz),  $\text{CDCl}_3$ ,  $\delta$ , 1.46 (s, 9H), 2.95 (m, 2H), 4.37 (m, 1H), 5.13 (s, 2H), 5.62 (m, 1H), 7.38 (m, 5H), 9.65 (s, 1H), MS(FAB+) 314.3 (M+Li).

### Step C

Preparation of 3-N-t-Boc-amino-4-hydroxy-4-phenyl-(3S)-butyric acid benzyl ester

To a diethyl ether (150 mL) solution of aldehyde (5.0 g, 15 mmole) prepared in Step B at  $-40^\circ\text{C}$  (acetonitrile/dry ice bath) was added in a dropwise fashion a 3.0 M solution of phenyl magnesium bromide in diethyl ether (10.8 mL, 32.6 mmole, 2 equivalents). The resulting mixture was stirred for 15 minutes and warmed to

- 626 -

room temperature. After several minutes the mixture was poured into 1 M  $K_2HPO_4$ . The aqueous layer was extracted again with ether, the combined ether layers washed with aqueous  $NaHCO_3$ , dried ( $Na_2SO_4$ ) and evaporated to give an oil (5.66 g) that was used in the next step without further purification:  $^1H$  NMR (300 MHz),  $CDCl_3$ ,  $\delta$ , 1.4 (multiple singlets, 9H), 2.65 (m, 2H), 4.18 (m, 1H), 5.15 (m, 2H), 7.4 (m, 10H); MS(FAB+) 392.4 (M+Li+).

10 Step D


Preparation of 2-phenyl-3-N-t-Boc-amino-5-oxo-3S-furan

The hydroxy-ester product of Step C (5.31 g, 13.8 mmole) was taken up in benzene (100 mL) a catalytic amount of camphor sulfonic acid was added and the solution refluxed (Dean-Stark) for five hours and the solvent removed. Conversion to lactone was 50% so the reaction was reconstituted and refluxed for a further 6 hours. Solvent was removed and the resulting oil taken up in ethyl acetate. The organic layer was washed with aqueous saturated  $NaHCO_3$ , dried ( $Na_2SO_4$ ) and evaporated to give a mixture of the desired diastereomeric lactones as a viscous oil in a 2:1 ratio and benzyl alcohol:  $^1H$  NMR (300 MHz),  $CDCl_3$ ,  $\delta$ , 1.35, 1.45 (s, 2:1, 9H), 2.75 (m, 2H), 4.5, 4.75 (m, 2:1, 1H), 4.7 (s, 2H), 5.1 (m, 1H), 5.7 (d, 1H), 7.35 (m, 10H); MS(FAB+) 284.6 (M+Li+).

- 627 -

Step E

Preparation of 2-phenyl-3-amino-5-oxo-3S-furane,  
hydrochloride




The lactone (0.94 g, 3.4 mmole) prepared in Step D was treated with 4 N HCl in dioxane (20 mL) at room temperature until gas evolution ceased. Excess HCl was removed by evaporation and the desired amino lactone isolated as a white crystalline solid that was dessicated (0.48 g, 66%):  $^1\text{H}$  NMR (300 MHz),  $d_6$  DMSO,  $\delta$ , 3.05 (m, 2H), 4.4 (m, 1H), 5.85 (d, 1H), 7.4 (s, 5H), 8.2 (bs, 3H); MS(FAB+) 178 (M+H+).

- 628 -


Step F

## Preparation f




The above compound is prepared using substantially the procedure of Example 374, substituting the product of Step E for the compound prepared in Example 233, Step B. The desired product is isolated by C-18 RPHPLC.

- 629 -


EXAMPLE 394

The above compound is prepared following substantially the procedure of Example 393, substituting 4-fluorophenyl magnesium bromide for phenyl magnesium bromide in Step C.


EXAMPLE 395

The above compound is prepared following substantially the procedure of Example 393, substituting 4-chlorophenyl magnesium bromide for phenyl magnesium bromide in Step C.

- 630 -


EXAMPLE 396

The above compound is prepared following substantially the procedure of Example 393, substituting 4-bromophenyl magnesium bromide for phenyl magnesium bromide in Step C.


EXAMPLE 397

The above compound is prepared following substantially the procedure of Example 393, substituting vinyl magnesium bromide for phenyl magnesium bromide in Step C.

- 631 -

EXAMPLE 398


10 The above compound is prepared following substantially the procedure of Example 393, substituting ethynylmagnesium bromide for phenyl magnesium bromide in Step C.

EXAMPLE 399

25 The above compound is prepared following substantially the procedure of Example 393, substituting allylmagnesium bromide for phenyl magnesium bromide in Step C.




- 632 -

EXAMPLE 400

10        The above compound is prepared following substantially the procedure of Example 393, substituting cyclopentylmagnesium bromide for phenyl magnesium bromide in Step C.


15        EXAMPLE 401



25        The above compound is prepared following substantially the procedure of Example 393, substituting phenylethynylmagnesium bromide for phenyl magnesium bromide in Step C.


30

- 633 -

EXAMPLE 402

10        The above compound is prepared following substantially the procedure of Example 393, substituting methylmagnesium bromide for phenyl magnesium bromide in Step C.


15        EXAMPLE 403




25        The above compound is prepared following substantially the procedure of Example 393, substituting isopropylmagnesium bromide for phenyl magnesium bromide in Step C.

30

- 634 -

EXAMPLE 404Step A

Preparation of 4-bromomagnesium-1,2-(methylenedioxy)benzene



20

To 1.74 gm (0.072 mole) freshly-ground magnesium in 100 mL dry THF in a 250 mL round bottom flask was added in a dropwise fashion 13.1 gm (0.062 mole) 4-bromo-1,2-(methylenedioxy)benzene in 50 mL dry THF. The reaction mixture was sonicated during the addition and the reaction temperature maintained below 50°C by use of a water bath. Upon completion of reaction the mixture was filtered and used in the next step.


- 635 -

Step B

Preparation of


5

10




The above compound is prepared following substantially the procedure of Example 393, substituting the grignard of Step A for phenyl magnesium bromide in Example 393, Step C.

- 636 -

EXAMPLE 405Step A

Preparation of




The above compound is prepared according to the procedure of Example 55, Step A, substituting methyl-2-formylbenzoate for 2-furancarboxaldehyde.

20

Step B

Preparation of




The above compound is prepared according to the procedure of Example 55, Steps B and C, substituting the product of Step A for the product of Example 55, Step A.

- 637 -


EXAMPLE 406

5

Step A

10 Preparation of

15




20

The product of Example 393, Step C is oxidized to the above ketone using the procedure of Example 393, Step B.

Step B

Preparation of

25




30

The above product is prepared using the procedure of Example 393, Step E using the product of Step A above.

- 638 -

Step C

## Preparation f



The Example compound is prepared using substantially the procedure of Example 374, substituting the product of Step B for the compound prepared in Example 233, Step B.

The desired product is obtained by converting the benzyl ester to the corresponding carboxylic acid by hydrolysis using substantially the procedure of Example 4 and isolating the desired product by C-18 RPHPLC.

- 639 -


Example 407-414

Using the procedure of Example 406, substituting the appropriate protected aspartyl alcohols prepared in  
5 Examples 394-403 for the aspartyl alcohol of Example 406, Step A, the following representative compounds are prepared:




- 640 -


Ex. 407




Ex. 408



Ex. 409




Ex. 410




- 641 -


Ex. 411




Ex. 412



Ex. 413




Ex. 414



- 642 -


Example 415

5



10

Preparation of




15

Step A

To the product of Example 23, Step A in DMF is added excess 1,3-diamino-2-hydroxypropane and catalytic DMAP and the solution heated until substantially complete conversion of the starting S-methylisothiuronium salt is achieved. The desired product may be isolated by precipitation of the zwitterion or by preparative C-18 RPHPLC (for a related procedure see U.S. Patent 2,899,426). After drying to remove water, the hydrochloride salt is formed by stirring the zwitterion in excess 4N HCl in dioxane (Aldrich) and isolating the HCl salt by filtration.

- 643 -




Step B

The above compound is prepared using substantially the procedure of Example 233, substituting the product of Step A for GHIA hydrochloride in Example 233, Step C.






- 644 -






Example 416-439

Using substantially the procedure of Example 415, substituting the appropriate amine for (RS)-4-amino-6-chloro-8-bromo-hydrocoumarin hydrochloride the following representative compounds may be prepared:








- 645 -









- 647 -









- 649 -



Ex. 438




Ex. 439



- 650 -

Example 440Step A

The above compound was prepared using the procedure of Example 233, Steps A and B, substituting 3,5-dichlorosalicylaldehyde for 3-bromo-5-chlorosalicylaldehyde in Step A. NMR and MS were consistent with the proposed structure (HCl salt).


Step B

The above compound is prepared by treating the product of Step A with dry HCl gas in methanol in a

- 651 -


suitable reactor while maintaining vigorous stirring. Upon completion of reaction excess HCl is removed under vacuum and the solution concentrated to dryness. The crude product is used in the next step. Alternatively, the product may be isolated by C-18 RPHPLC and lyophilized to obtain substantially pure material.

### Step C




The above compound is prepared by taking the product of Step B and dissolving in DMF. To the stirred solution is added an equimolar amount of both di-tert butyl dicarbonate and triethylamine with a catalytic amount of DMAP. Upon completion of the reaction volatiles are removed under vacuum and the product partitioned between dilute aqueous hydrochloric acid and ethyl acetate. The organic layer is washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to provide substantially the above compound that may be employed in the next step without further purification. Alternatively, the product may be isolated by C-18 RPHPLC and lyophilized to obtain substantially pure material.

- 652 -

Step D


The above compound is prepared by adding under an inert atmosphere an equivalent of acetic anhydride or acetyl chloride and an equivalent of triethylamine to a stirred solution of the product from Step C in DMF. Upon completion of reaction volatiles are removed under vacuum and the reaction residue partitioned between dilute aqueous hydrochloric acid and ethyl acetate. The organic layer is washed with aqueous sodium bicarbonate, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to provide substantially the above compound that may be employed in the next step without further purification. Alternatively, the product may be isolated by C-18 RPHPLC and lyophilized to obtain substantially pure material.

Step E

- 653 -


The above compound is prepared by treating the product of Step D with 4 N HCl in di xane with vigorous stirring. Shortly after cessation of gas evolution excess HCl gas is removed *in vacuo* and the reaction mixture concentrated at less than about 40°C. The product is trituated with diethyl ether to obtain substantially the desired product. Alternatively, the product may be isolated by C-18 RPHPLC and lyophilized to obtain substantially pure mater.

#### Step F




The above compound is prepared according to the procedure of Example 230, Step B, substituting the product of Step E for the product of Example 230, Step A.

- 654 -

Example 441

The above compound is prepared by treating a DMF mixture of the compound of Example 225 with two equivalents of N-methylmorpholine and one equivalent of acetic anhydride or acetyl chloride. Upon completion of the reaction the desired product may be isolated by C-18 RPHPLC and lyophilization.





Example 442

The above compound is prepared by treating a DMF mixture of the compound of Example 225 with two equivalents of N-methylmorpholine and one equivalent of benzoic anhydride or benzoyl chloride. Upon completion of the reaction the desired product may be isolated by C-18 RPHPLC and lyophilization.

- 655 -

Example 443-452


Using substantially the procedure of Example 230, Step B and substituting the appropriate amine for the product of Example 230, Step A, the following  
5 representative compounds may be prepared:






- 656 -


Ex. 447



Ex. 448




Ex. 449




- 657 -


Ex. 450



Ex. 451



Ex. 452




- 658 -


Example 453-460

Using the procedure of Example 393, substituting the appropriate amine hydrochloride for the product of Step E in Step F and substituting GIHA HCl for 3-guanidino-5-trifluoromethylhippuric acid in Step F the following representative compounds may be prepared:

Ex. 453




Ex. 454




- 659 -


Ex. 455




Ex. 456




Ex. 457




Ex. 458



Ex. 459



Ex. 460




- 661 -


Example 461

Using the procedure of Example 406, substituting the appropriate protected aspartyl alcohol prepared in Examples 394-403 for the aspartyl alcohol of Example 406, Step A, and substituting GIHA HCl for 3-guanidino-5-trifluoromethylhippuric acid in Example 393, Step F the following representative compounds may be prepared:


Ex. 461



Ex. 462




Ex. 463




- 662 -


Ex. 464



Ex. 465




Ex. 466




- 663 -


Ex. 467



Ex. 468



Ex. 469






- 664 -

Example 470

## Preparation of

Step A

To 3,4,5,6-tetrahydro-2-pyrimidinethiol (Aldrich) (5.0 g, 0.043 mole) and triethylamine (8.7 g, 0.086 mole) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added dropwise and at ice bath temperature, phenylchloroformate [(Aldrich) 13.5 g, 0.086 mole)]. The reaction was then stirred overnight at room temperature. The precipitate was filtered and washed with CH<sub>2</sub>Cl<sub>2</sub>. The CH<sub>2</sub>Cl<sub>2</sub> filtrate was washed with H<sub>2</sub>O (3X), dried over MgSO<sub>4</sub> and removed under vacuum. The residue was recrystallized from 50% EtOAc/Hexane to yield 9.03 g of 3,4,5,6-tetrahydro-2-pyrimidinethione-N,N'-diphenylcarbamate as a yellow solid.

MS and NMR are consistent with the desired structure.

Step B

To the product from Example 282, Step C (200 mg, 0.00042 mole), the product from Step A above (150 mg, 0.00042 mole) and triethylamine (142 mg, 0.0014 mole) in 3 mL DMF was added 250 mg (0.00046 mole) HgCl<sub>2</sub> at ice bath temperature. The reaction was stirred at ice bath temperature for ½ hour and at room temperature for 2 hours. 100 mg additional HgCl<sub>2</sub> was added and the reaction was stirred overnight at 60°C. Excess ethyl acetate was added and the slurry was filtered through

- 665 -


celite. The filtrate was washed with H<sub>2</sub>O (3X), passed through a pad of silica gel and the product isolated by silica gel chromatography to yield the above compound (110 mg) as a white solid.

5. MS and NMR were consistent with the desired structure.

- 666 -

Example 471

## Preparation of

Step A

15 To 3,4,5,6-tetrahydro-2-pyrimidinethiol (Aldrich) (10 g, 0.086 mole) in absolute ethanol (75 mL) is added methyl iodide (12.2 g, 0.086 mole). The reaction was stirred at reflux for 2.5 hours. The solvent was removed under vacuum and the residue dried to yield

20 3,4,5,6-tetrahydro-2-methylthiopyrimidine·HI (22 g) as a white solid.

MS and NMR were consistent with the desired structure.

25 Step B

To the product from Step A above (5.16 g, 0.02 mole) and triethylamine (4.1 g, 0.04 mole) in  $\text{CH}_2\text{Cl}_2$  (25 mL) was added phenylchloroformate (Aldrich) (3.13 g, 0.02 mole) dropwise at ice bath temperature. The

30 reaction was then stirred overnight at room temperature. The precipitate was filtered and washed with  $\text{CH}_2\text{Cl}_2$ . The  $\text{CH}_2\text{Cl}_2$  from the filtrate was washed with  $\text{H}_2\text{O}$  (3X), dried over  $\text{MgSO}_4$  and removed under vacuum to yield 3,4,5,6-tetrahydro-2-methylthiopyrimidine-N-

35 phenylcarbamate (4.8 g) as a white solid.

MS and NMR were consistent with the desired structure.

- 667 -

Step C

To the product from Step B above (2 g, 0.008 mole) in CH<sub>2</sub>CN (12 mL) was added the product of Example M, Step B (1.84 g, 0.008 mole). The reaction was stirred at reflux overnight and the product isolated by RPHPLC to yield 3,4,5,6-tetrahydro-N-phenylcarbamyl-2-pyrimidine-m-aminohippuric acid·TFA (1 g) as a white solid.

10 Step D


The above compound was prepared according to the methodology of Example 174, substituting an equivalent amount of 3,5-dichlorobenzaldehyde for 3,4-dichlorobenzaldehyde in Example 174, Step A and substituting an equivalent amount of the product from Step C above for m-guanidinohippuric acid·HCl in Example 174, Step B.

MS and NMR were consistent with the desired structure.

- 668 -

Example 472

## Preparation of



**Step A**

To 2-methylthio-2-imidazoline·HI (Aldrich) (10 g, 0.041 mole) and triethylamine (4.14 g, 0.041 mole) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added BOC anhydride (Aldrich) (8.94 g, 0.041 mole) at ice bath temperature. The reaction was stirred overnight at room temperature. The CH<sub>2</sub>Cl<sub>2</sub> was washed with H<sub>2</sub>O (3 X), dried over MgSO<sub>4</sub>, washed with H<sub>2</sub>O (3 X), dried over MgSO<sub>4</sub> and removed under vacuum to yield N-BOC-2-methylthio-2-imidazoline (8.1 g) as a clear liquid which turned to a white solid upon standing.

MS and NMR were consistent with the desired structure.

**25     Step B**

To the product of Step A above (2.7 g, 0.0124 mole) in  $\text{CH}_3\text{CN}$  (6 mL) was added 3-amino-5-trifluoromethylbenzoic acid (synthesized by catalytic hydrogenation (Pd/C) of 3-nitro-5-trifluorobenzoic acid (Lancaster) followed by treatment with HCl) (3 g, 0.0124 mole). The reaction was stirred at 35-40°C for 10 days. After cooling to room temperature, the precipitate was filtered, washed with  $\text{CH}_3\text{CN}$  and dried to yield 3-(N-BOC-4,5-dihydroimidazol-2-yl)amino-5-trifluoromethylbenzoic acid HCl (3.2 g) as a white solid.

- 669 -

MS and NMR were consistent with the desired structure.

Step C


- 5       The above compound was prepared according to the methodology of Example 200, substituting an equivalent amount of the product from Step B above for the product from Step A in Example 199, Step B and by additionally treating the intermediate ethyl ester, N-BOC derivative
- 10       with TFA for 1 hour to remove the BOC protecting group.

MS and NMR were consistent with the desired structure.

- 670 -

Example 473

## Preparation of

Step A

To 3-amino-5-trifluoromethylhippuric acid hydrochloride [prepared according to Example M, Steps A and B substituting 3-nitro-5-trifluoromethylbenzoyl chloride (prepared from 3-nitro-5-trifluoromethylbenzoic acid (Lancaster) and thionyl chloride for *m*-nitrobenzoyl chloride in Example M, Step A] (3 g, 0.01 mole) in CH<sub>3</sub>CN (5 mL) was added the product from Example 472, Step A (2.2 g, 0.01 mole). The reaction was stirred at 35°C for 3 days then at reflux for 4 hours. After cooling, the CH<sub>3</sub>CN was decanted off, the residue slurried several times in ether (ether decanted off) and then dried to yield 3-((4,5-dihydro-1H-imidazol-2-yl)amino)-5-trifluoromethylhippuric acid·HCl (2.5 g) as a white solid.

MS and NMR were consistent with the desired structure.

Step B


The above compound was prepared according to the methodology of Example 210, substituting an equivalent amount of the product from Step A above for *m*-guanidinohippuric acid·HCl in Example 174, Step B.

MS and NMR were consistent with the desired structure.

- 671 -

Example 474

## Preparation of

Step A

To 2-methylthio-2-imidazoline·HI (Aldrich) (10 g, 0.041 mole) and triethylamine (8.3 g, 0.0082 mole) in  $\text{CH}_2\text{Cl}_2$  (50 mL) was added ethylchloroformate (Aldrich) (4.5 g, 0.041 mole) dropwise at ice bath temperature. The reaction was stirred overnight at room temperature. The precipitate was filtered and washed with  $\text{CH}_2\text{Cl}_2$ . The  $\text{CH}_2\text{Cl}_2$  from the filtrate was washed with  $\text{H}_2\text{O}$  (3X), dried over  $\text{MgSO}_4$  and removed under vacuum to yield 2-methylthio-2-imidazoline-N-ethylcarbamate (7.1 g) as a clear yellow oil.

MS and NMR were consistent with the desired structure.

Step B

To the product from Step A above (5.73 g, 0.0305 mole) in  $\text{CH}_3\text{CN}$  (12 mL) was added m-aminohippuric acid·HCl (Example M, Step B) (7.02 g, 0.0305 mole). The reaction was stirred overnight at room temperature then at  $50^\circ\text{C}$  for 6 hours and at  $80^\circ\text{C}$  for 2 hours. After cooling to room temperature and stirring at room temperature overnight, the precipitate was filtered, washed with  $\text{CH}_3\text{CN}$  and dried to yield 3-(4,5-dihydro-N-ethylcarbamate-imidazol-2-yl)aminohippuric acid·HCl (9.6 g) as a white solid.

MS and NMR were consistent with the desired structure.



- 672 -

Step C

The above compound was prepared according to the methodology of Example 174, substituting an equivalent amount of the product from Step B above for


- 5 m-aminohippuric acid in Example 174, Step B and an equivalent amount of the product from Example 230, Step A for the product from Example 174, Step A in Example 174, Step B.

- 10 MS and NMR were consistent with the desired structure.

- 673 -

Example 475

## Preparation of




10

The above compound was prepared according to the methodology of Example 474, substituting an equivalent amount of phenylchloroformate (Aldrich) for ethylchloroformate in Example 474, Step A and by heating the reaction mixture at 70°C for 8 hours then room temperature for 2 days in Example 474, Step B.





MS and NMR were consistent with the desired structure.

- 674 -





Using the methodologies, reagents and conditions exemplified in the schemes and examples of this disclosure (or the synthesis of reagents from readily available starting materials via methodologies known to those skilled in the art), the following compounds of the present invention are synthesized:

Examples 476-517





- 675 -








- 676 -







- 677 -















- 679 -














- 681 -








- 684 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	518	OH	Cl	Cl	Br	H	H
	519	OH	Cl	Cl	OH	H	H
	520	OH	Cl	Cl	NO <sub>2</sub>	H	H
	521	OH	Cl	Cl	I	H	H
	522	OH	Cl	Cl	Cl	H	H
15	523	OH	Cl	Cl	Cl	H	Cl
	524	OH	Cl	Cl	OMe	H	H
	525	OH	Cl	Cl	H	CF <sub>3</sub>	H
	526	OH	Cl	Cl	H	OH	H
	527	OH	Cl	Cl	H	OMe	H
20	528	OH	Cl	Cl	H	Cl	H
	529	OH	Cl	Cl	H	Cl	Cl
	530	OH	Cl	Cl	H	Br	H
	531	OH	Cl	Cl	Cl	OH	H
	532	OH	Cl	Cl	Br	OH	H
25	533	OH	Cl	Cl	I	OH	H

- 685 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	534	OH	Br	Cl	Br	H	H
	535	OH	Br	Cl	OH	H	H
	536	OH	Br	Cl	NO <sub>2</sub>	H	H
	537	OH	Br	Cl	I	H	H
	538	OH	Br	Cl	Cl	H	H
15	539	OH	Br	Cl	Cl	H	Cl
	540	OH	Br	Cl	OMe	H	H
	541	OH	Br	Cl	H	CF <sub>3</sub>	H
	542	OH	Br	Cl	H	OH	H
	543	OH	Br	Cl	H	OMe	H
20	544	OH	Br	Cl	H	Cl	H
	545	OH	Br	Cl	H	Cl	Cl
	546	OH	Br	Cl	H	Br	H
	547	OH	Br	Cl	Cl	OH	H
	548	OH	Br	Cl	Br	OH	H
25	549	OH	Br	Cl	I	OH	H

- 686 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	550	OH	I	Cl	Br	H	H
	551	OH	I	Cl	OH	H	H
	552	OH	I	Cl	NO <sub>2</sub>	H	H
	553	OH	I	Cl	I	H	H
	554	OH	I	Cl	Cl	H	H
15	555	OH	I	Cl	Cl	H	Cl
	556	OH	I	Cl	OMe	H	H
	557	OH	I	Cl	H	CF <sub>3</sub>	H
	558	OH	I	Cl	H	OH	H
	559	OH	I	Cl	H	OMe	H
20	560	OH	I	Cl	H	Cl	H
	561	OH	I	Cl	H	Cl	Cl
	562	OH	I	Cl	H	Br	H
	563	OH	I	Cl	CF <sub>3</sub>	H	H
	564	OH	I	Cl	Cl	OH	H
25	565	OH	I	Cl	Br	OH	H
	566	OH	I	Cl	I	OH	H

- 687 -

5




	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	567	H	Br	Cl	Br	H	H
	568	H	Br	Cl	OH	H	H
	569	H	Br	Cl	NO <sub>2</sub>	H	H
	570	H	Br	Cl	I	H	H
	571	H	Br	Cl	Cl	H	H
15	572	H	Br	Cl	Cl	H	Cl
	573	H	Br	Cl	OMe	H	H
	574	H	Br	Cl	H	CF <sub>3</sub>	H
	575	H	Br	Cl	H	OH	H
	576	H	Br	Cl	H	OMe	H
20	577	H	Br	Cl	H	Cl	H
	578	H	Br	Cl	H	Cl	Cl
	579	H	Br	Cl	H	Br	H
	580	H	Br	Cl	Cl	OH	H
	581	H	Br	Cl	Br	OH	H
25	582	H	Br	Cl	I	OH	H



- 688 -


5




	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	583	H	Br	Br	Br	H	H
	584	H	Br	Br	OH	H	H
	585	H	Br	Br	NO <sub>2</sub>	H	H
	586	H	Br	Br	I	H	H
	587	H	Br	Br	Cl	H	H
15	588	H	Br	Br	Cl	H	Cl
	589	H	Br	Br	OMe	H	H
	590	H	Br	Br	H	CF <sub>3</sub>	H
	591	H	Br	Br	H	OH	H
	592	H	Br	Br	H	OMe	H
20	593	H	Br	Br	H	Cl	H
	594	H	Br	Br	H	Cl	Cl
	595	H	Br	Br	H	Br	H
	596	H	Br	Br	Cl	OH	H
	597	H	Br	Br	Br	OH	H
25	598	H	Br	Br	I	OH	H

- 689 -

5




	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	599	H	Br	I	Br	H	H
	600	H	Br	I	OH	H	H
	601	H	Br	I	NO <sub>2</sub>	H	H
	602	H	Br	I	I	H	H
	603	H	Br	I	Cl	H	H
15	604	H	Br	I	Cl	H	Cl
	605	H	Br	I	OMe	H	H
	606	H	Br	I	H	CF <sub>3</sub>	H
	607	H	Br	I	H	OH	H
	608	H	Br	I	H	OMe	H
20	609	H	Br	I	H	Cl	H
	610	H	Br	I	H	Cl	Cl
	611	H	Br	I	H	Br	H
	612	H	Br	I	Cl	OH	H
	613	H	Br	I	Br	OH	H
25	614	H	Br	I	I	OH	H



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	615	H	I	I	Br	H	H
	616	H	I	I	OH	H	H
	617	H	I	I	NO <sub>2</sub>	H	H
	618	H	I	I	I	H	H
	619	H	I	I	Cl	H	H
15	620	H	I	I	Cl	H	Cl
	621	H	I	I	OMe	H	H
	622	H	I	I	H	CF <sub>3</sub>	H
	623	H	I	I	H	OH	H
	624	H	I	I	H	OMe	H
20	625	H	I	I	H	Cl	H
	626	H	I	I	H	Cl	Cl
	627	H	I	I	H	Br	H
	628	H	I	I	Cl	OH	H
	629	H	I	I	Br	OH	H
25	630	H	I	I	I	OH	H


- 691 -

5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	631	H	Cl	Cl	Br	H	H
	632	H	Cl	Cl	OH	H	H
	633	H	Cl	Cl	NO <sub>2</sub>	H	H
	634	H	Cl	Cl	I	H	H
	635	H	Cl	Cl	Cl	H	H
15	636	H	Cl	Cl	Cl	H	Cl
	637	H	Cl	Cl	OMe	H	H
	638	H	Cl	Cl	H	CF <sub>3</sub>	H
	639	H	Cl	Cl	H	OH	H
	640	H	Cl	Cl	H	OMe	H
20	641	H	Cl	Cl	H	Cl	H
	642	H	Cl	Cl	H	Cl	Cl
	643	H	Cl	Cl	H	Br	H
	644	H	Cl	Cl	Cl	OH	H
	645	H	Cl	Cl	Br	OH	H
25	646	H	Cl	Cl	I	OH	H


- 692 -



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	647	OH	Cl	Cl	Br	H	H
	648	OH	Cl	Cl	OH	H	H
	649	OH	Cl	Cl	NO <sub>2</sub>	H	H
	650	OH	Cl	Cl	I	H	H
	651	OH	Cl	Cl	Cl	H	H
15	652	OH	Cl	Cl	Cl	H	Cl
	653	OH	Cl	Cl	OMe	H	H
	654	OH	Cl	Cl	H	CF <sub>3</sub>	H
	655	OH	Cl	Cl	H	OH	H
	656	OH	Cl	Cl	H	OMe	H
20	657	OH	Cl	Cl	H	Cl	H
	658	OH	Cl	Cl	H	Cl	Cl
	659	OH	Cl	Cl	H	Br	H
	660	OH	Cl	Cl	Cl	OH	H
	661	OH	Cl	Cl	Br	OH	H
25	662	OH	Cl	Cl	I	OH	H


- 693 -

5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	663	OH	Br	Cl	Br	H	H
	664	OH	Br	Cl	OH	H	H
	665	OH	Br	Cl	NO <sub>2</sub>	H	H
	666	OH	Br	Cl	I	H	H
	667	OH	Br	Cl	Cl	H	H
15	668	OH	Br	Cl	Cl	H	Cl
	669	OH	Br	Cl	OMe	H	H
	670	OH	Br	Cl	H	CF <sub>3</sub>	H
	671	OH	Br	Cl	H	OH	H
	672	OH	Br	Cl	H	OMe	H
20	673	OH	Br	Cl	H	Cl	H
	674	OH	Br	Cl	H	Cl	Cl
	675	OH	Br	Cl	H	Br	H
	676	OH	Br	Cl	Cl	OH	H
	677	OH	Br	Cl	Br	OH	H
25	678	OH	Br	Cl	I	OH	H


- 694 -



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	679	OH	I	Cl	Br	H	H
	680	OH	I	Cl	OH	H	H
	681	OH	I	Cl	NO <sub>2</sub>	H	H
	682	OH	I	Cl	I	H	H
	683	OH	I	Cl	Cl	H	H
15	684	OH	I	Cl	Cl	H	Cl
	685	OH	I	Cl	OMe	H	H
	686	OH	I	Cl	H	CF <sub>3</sub>	H
	687	OH	I	Cl	H	OH	H
	688	OH	I	Cl	H	OMe	H
20	689	OH	I	Cl	H	Cl	H
	690	OH	I	Cl	H	Cl	Cl
	691	OH	I	Cl	H	Br	H
	692	OH	I	Cl	Cl	OH	H
	693	OH	I	Cl	Br	OH	H
25	694	OH	I	Cl	I	OH	H

- 695 -

5




	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	695	H	Cl	Cl	Br	H	H
	696	H	Cl	Cl	OH	H	H
	697	H	Cl	Cl	NO <sub>2</sub>	H	H
	698	H	Cl	Cl	I	H	H
	699	H	Cl	Cl	Cl	H	H
15	700	H	Cl	Cl	Cl	H	Cl
	701	H	Cl	Cl	OMe	H	H
	702	H	Cl	Cl	H	CF <sub>3</sub>	H
	703	H	Cl	Cl	H	OH	H
	704	H	Cl	Cl	H	OMe	H
20	705	H	Cl	Cl	H	Cl	H
	706	H	Cl	Cl	H	Cl	Cl
	707	H	Cl	Cl	H	Br	H
	708	H	Cl	Cl	Cl	OH	H
	709	H	Cl	Cl	Br	OH	H
25	710	H	Cl	Cl	I	OH	H



- 696 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	711	H	Br	Cl	Br	H	H
	712	H	Br	Cl	OH	H	H
	713	H	Br	Cl	NO <sub>2</sub>	H	H
	714	H	Br	Cl	I	H	H
	715	H	Br	Cl	Cl	H	H
15	716	H	Br	Cl	Cl	H	Cl
	717	H	Br	Cl	OMe	H	H
	718	H	Br	Cl	H	CF <sub>3</sub>	H
	719	H	Br	Cl	H	OH	H
	720	H	Br	Cl	H	OMe	H
20	721	H	Br	Cl	H	Cl	H
	722	H	Br	Cl	H	Cl	Cl
	723	H	Br	Cl	H	Br	H
	724	H	Br	Cl	Cl	OH	H
	725	H	Br	Cl	Br	OH	H
	726	H	Br	Cl	I	OH	H

25


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	727	H	Br	Br	Br	H	H
	728	H	Br	Br	OH	H	H
	729	H	Br	Br	NO <sub>2</sub>	H	H
	730	H	Br	Br	I	H	H
	731	H	Br	Br	Cl	H	H
15	732	H	Br	Br	Cl	H	Cl
	733	H	Br	Br	OMe	H	H
	734	H	Br	Br	H	CF <sub>3</sub>	H
	735	H	Br	Br	H	OH	H
	736	H	Br	Br	H	OMe	H
20	737	H	Br	Br	H	Cl	H
	738	H	Br	Br	H	Cl	Cl
	739	H	Br	Br	H	Br	H
	740	H	Br	Br	Cl	OH	H
	741	H	Br	Br	Br	OH	H
25	742	H	Br	Br	I	OH	H

- 698 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	743	H	Br	I	Br	H	H
	744	H	Br	I	OH	H	H
	745	H	Br	I	NO <sub>2</sub>	H	H
	746	H	Br	I	I	H	H
	747	H	Br	I	Cl	H	H
15	748	H	Br	I	Cl	H	Cl
	749	H	Br	I	OMe	H	H
	750	H	Br	I	H	CF <sub>3</sub>	H
	751	H	Br	I	H	OH	H
	752	H	Br	I	H	OMe	H
20	753	H	Br	I	H	Cl	H
	754	H	Br	I	H	Cl	Cl
	755	H	Br	I	H	Br	H
	756	H	Br	I	Cl	OH	H
	757	H	Br	I	Br	OH	H
25	758	H	Br	I	I	OH	H

- 699 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	759	H	I	I	Br	H	H
	760	H	I	I	OH	H	H
	761	H	I	I	NO <sub>2</sub>	H	H
	762	H	I	I	I	H	H
	763	H	I	I	Cl	H	H
15	764	H	I	I	Cl	H	Cl
	765	H	I	I	OMe	H	H
	766	H	I	I	H	CF <sub>3</sub>	H
	767	H	I	I	H	OH	H
	768	H	I	I	H	OMe	H
20	769	H	I	I	H	Cl	H
	770	H	I	I	H	Cl	Cl
	771	H	I	I	H	Br	H
	772	H	I	I	Cl	OH	H
	773	H	I	I	Br	OH	H
25	774	H	I	I	I	OH	H


- 700 -

5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	775	OH	Cl	Cl	Br	H	H
	776	OH	Cl	Cl	OH	H	H
	777	OH	Cl	Cl	NO <sub>2</sub>	H	H
	778	OH	Cl	Cl	I	H	H
	779	OH	Cl	Cl	Cl	H	H
15	780	OH	Cl	Cl	Cl	H	Cl
	781	OH	Cl	Cl	OMe	H	H
	782	OH	Cl	Cl	H	CF <sub>3</sub>	H
	783	OH	Cl	Cl	H	OH	H
	784	OH	Cl	Cl	H	OMe	H
20	785	OH	Cl	Cl	H	Cl	H
	786	OH	Cl	Cl	H	Cl	Cl
	787	OH	Cl	Cl	H	Br	H
	788	OH	Cl	Cl	Cl	OH	H
	789	OH	Cl	Cl	Br	OH	H
25	790	OH	Cl	Cl	I	OH	H

- 701 -




<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	791	OH	Br	Cl	Br	H	H
	792	OH	Br	Cl	OH	H	H
	793	OH	Br	Cl	NO <sub>2</sub>	H	H
	794	OH	Br	Cl	I	H	H
	795	OH	Br	Cl	Cl	H	H
	796	OH	Br	Cl	Cl	H	Cl
15	797	OH	Br	Cl	OMe	H	H
	798	OH	Br	Cl	H	CF <sub>3</sub>	H
	799	OH	Br	Cl	H	OH	H
	800	OH	Br	Cl	H	OMe	H
	801	OH	Br	Cl	H	Cl	H
	802	OH	Br	Cl	H	Cl	Cl
20	803	OH	Br	Cl	H	Br	H
	804	OH	Br	Cl	Cl	OH	H
	805	OH	Br	Cl	Br	OH	H
	806	OH	Br	Cl	I	OH	H

25

- 702 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	807	OH	I	Cl	Br	H	H
	808	OH	I	Cl	OH	H	H
	809	OH	I	Cl	NO <sub>2</sub>	H	H
	810	OH	I	Cl	I	H	H
	811	OH	I	Cl	Cl	H	H
15	812	OH	I	Cl	Cl	H	Cl
	813	OH	I	Cl	OMe	H	H
	814	OH	I	Cl	H	CF <sub>3</sub>	H
	815	OH	I	Cl	H	OH	H
	816	OH	I	Cl	H	OMe	H
20	817	OH	I	Cl	H	Cl	H
	818	OH	I	Cl	H	Cl	Cl
	819	OH	I	Cl	H	Br	H
	820	OH	I	Cl	Cl	OH	H
	821	OH	I	Cl	Br	OH	H
25	822	OH	I	Cl	I	OH	H

- 703 -

5




<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	823	H	Cl	Cl	Br	H	H
	824	H	Cl	Cl	H	H	H
	825	H	Cl	Cl	NO <sub>2</sub>	H	H
	826	H	Cl	Cl	I	H	H
	827	H	Cl	Cl	Cl	H	H
15	828	H	Cl	Cl	Cl	H	Cl
	829	H	Cl	Cl	OMe	H	H
	830	H	Cl	Cl	H	CF <sub>3</sub>	H
	831	H	Cl	Cl	H	OH	H
	832	H	Cl	Cl	H	OMe	H
20	833	H	Cl	Cl	H	Cl	H
	834	H	Cl	Cl	H	Cl	Cl
	835	H	Cl	Cl	H	Br	H
	836	H	Cl	Cl	Cl	OH	H
	837	H	Cl	Cl	Br	OH	H
25	838	H	Cl	Cl	I	OH	H



- 704 -

5




<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	839	H	Br	Cl	Br	H	H
	840	H	Br	Cl	OH	H	H
	841	H	Br	Cl	NO <sub>2</sub>	H	H
	842	H	Br	Cl	I	H	H
	843	H	Br	Cl	Cl	H	H
15	845	H	Br	Cl	Cl	H	Cl
	846	H	Br	Cl	OMe	H	H
	847	H	Br	Cl	H	CF <sub>3</sub>	H
	848	H	Br	Cl	H	OH	H
	849	H	Br	Cl	H	OMe	H
20	850	H	Br	Cl	H	Cl	H
	851	H	Br	Cl	H	Cl	Cl
	852	H	Br	Cl	H	Br	H
	853	H	Br	Cl	Cl	OH	H
	854	H	Br	Cl	Br	OH	H
	855	H	Br	Cl	I	OH	H

25

- 705 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	856	H	Br	Br	Br	H	H
	857	H	Br	Br	OH	H	H
	858	H	Br	Br	NO <sub>2</sub>	H	H
	859	H	Br	Br	I	H	H
	860	H	Br	Br	Cl	H	H
15	861	H	Br	Br	Cl	H	Cl
	862	H	Br	Br	OMe	H	H
	863	H	Br	Br	H	CF <sub>3</sub>	H
	864	H	Br	Br	H	OH	H
	865	H	Br	Br	H	OMe	H
20	866	H	Br	Br	H	Cl	H
	867	H	Br	Br	H	Cl	Cl
	868	H	Br	Br	H	Br	H
	869	H	Br	Br	Cl	OH	H
	870	H	Br	Br	Br	OH	H
25	871	H	Br	Br	I	OH	H

- 706 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	872	H	Br	I	Br	H	H
	873	H	Br	I	OH	H	H
	874	H	Br	I	NO <sub>2</sub>	H	H
	875	H	Br	I	I	H	H
	876	H	Br	I	Cl	H	H
15	877	H	Br	I	Cl	H	Cl
	878	H	Br	I	OMe	H	H
	879	H	Br	I	H	CF <sub>3</sub>	H
	880	H	Br	I	H	OH	H
	881	H	Br	I	H	OMe	H
20	882	H	Br	I	H	Cl	H
	883	H	Br	I	H	Cl	Cl
	884	H	Br	I	H	Br	H
	885	H	Br	I	Cl	OH	H
	886	H	Br	I	Br	OH	H
25	887	H	Br	I	I	OH	H

- 707 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	888	H	I	I	Br	H	H
	889	H	I	I	OH	H	H
	890	H	I	I	NO <sub>2</sub>	H	H
	891	H	I	I	I	H	H
	892	H	I	I	Cl	H	H
15	893	H	I	I	Cl	H	Cl
	894	H	I	I	OMe	H	H
	895	H	I	I	H	CF <sub>3</sub>	H
	896	H	I	I	H	OH	H
	897	H	I	I	H	OMe	H
20	898	H	I	I	H	Cl	H
	899	H	I	I	H	Cl	Cl
	900	H	I	I	H	Br	H
	901	H	I	I	Cl	OH	H
	902	H	I	I	Br	OH	H
25	903	H	I	I	I	OH	H

- 708 -


5



<u>Example #</u>		<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	904	H	CF <sub>3</sub>	Br	Br	H	H
	905	H	CF <sub>3</sub>	Br	OH	H	H
	906	H	CF <sub>3</sub>	Br	NO <sub>2</sub>	H	H
	907	H	CF <sub>3</sub>	Br	I	H	H
	908	H	CF <sub>3</sub>	Br	Cl	H	H
15	909	H	CF <sub>3</sub>	Br	Cl	H	Cl
	910	H	CF <sub>3</sub>	Br	OMe	H	H
	911	H	CF <sub>3</sub>	Br	H	CF <sub>3</sub>	H
	912	H	CF <sub>3</sub>	Br	H	OH	H
	913	H	CF <sub>3</sub>	Br	H	OMe	H
20	914	H	CF <sub>3</sub>	Br	H	Cl	H
	915	H	CF <sub>3</sub>	Br	H	Cl	Cl
	916	H	CF <sub>3</sub>	Br	H	Br	H
	917	H	CF <sub>3</sub>	Br	CF <sub>3</sub>	H	H
	918	H	CF <sub>3</sub>	Br	H	H	H
25	919	H	CF <sub>3</sub>	Br	Cl	OH	H
	920	H	CF <sub>3</sub>	Br	Br	OH	H
	921	H	CF <sub>3</sub>	Br	I	OH	H

- 709 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	922	H	CF <sub>3</sub>	Br	Br	H	H
	923	H	CF <sub>3</sub>	Br	OH	H	H
	924	H	CF <sub>3</sub>	Br	NO <sub>2</sub>	H	H
	925	H	CF <sub>3</sub>	Br	I	H	H
	926	H	CF <sub>3</sub>	Br	Cl	H	H
	927	H	CF <sub>3</sub>	Br	Cl	H	Cl
15	928	H	CF <sub>3</sub>	Br	OMe	H	H
	929	H	CF <sub>3</sub>	Br	H	CF <sub>3</sub>	H
	930	H	CF <sub>3</sub>	Br	H	OH	H
	931	H	CF <sub>3</sub>	Br	H	OMe	H
	932	H	CF <sub>3</sub>	Br	H	Cl	H
	933	H	CF <sub>3</sub>	Br	H	Cl	Cl
20	934	H	CF <sub>3</sub>	Br	H	Br	H
	935	H	CF <sub>3</sub>	Br	CF <sub>3</sub>	H	H
	936	H	CF <sub>3</sub>	Br	H	H	H
	937	H	CF <sub>3</sub>	Br	Cl	OR	H
	938	H	CF <sub>3</sub>	Br	Br	OH	H
	939	H	CF <sub>3</sub>	Br	I	OH	H

- 710 -


5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	940	H	CF <sub>3</sub>	Br	Br	H	H
	941	H	CF <sub>3</sub>	Br	OH	H	H
	942	H	CF <sub>3</sub>	Br	NO <sub>2</sub>	H	H
	943	H	CF <sub>3</sub>	Br	I	H	H
	944	H	CF <sub>3</sub>	Br	Cl	H	H
15	945	H	CF <sub>3</sub>	Br	Cl	H	Cl
	946	H	CF <sub>3</sub>	Br	OMe	H	H
	947	H	CF <sub>3</sub>	Br	H	CF <sub>3</sub>	H
	948	H	CF <sub>3</sub>	Br	H	OH	H
	949	H	CF <sub>3</sub>	Br	H	OMe	H
20	950	H	CF <sub>3</sub>	Br	H	Cl	H
	951	H	CF <sub>3</sub>	Br	H	Cl	Cl
	952	H	CF <sub>3</sub>	Br	H	Br	H
	953	H	CF <sub>3</sub>	Br	CF <sub>3</sub>	H	H
	954	H	CF <sub>3</sub>	Br	H	H	H
25	955	H	CF <sub>3</sub>	Br	Cl	OH	H
	956	H	CF <sub>3</sub>	Br	Br	OH	H
	957	H	CF <sub>3</sub>	Br	I	OH	H

- 711 -

5




	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	958	H	CF <sub>3</sub>	I	Br	H	H
	959	H	CF <sub>3</sub>	I	OH	H	H
	960	H	CF <sub>3</sub>	I	NO <sub>2</sub>	H	H
	961	H	CF <sub>3</sub>	I	I	H	H
	962	H	CF <sub>3</sub>	I	Cl	H	H
	963	H	CF <sub>3</sub>	I	Cl	H	Cl
15	964	H	CF <sub>3</sub>	I	OMe	H	H
	965	H	CF <sub>3</sub>	I	H	CF <sub>3</sub>	H
	966	H	CF <sub>3</sub>	I	H	OH	H
	967	H	CF <sub>3</sub>	I	H	OMe	H
	968	H	CF <sub>3</sub>	I	H	Cl	H
20	969	H	CF <sub>3</sub>	I	H	Cl	Cl
	970	H	CF <sub>3</sub>	I	H	Br	H
	971	H	CF <sub>3</sub>	I	CF <sub>3</sub>	H	H
	972	H	CF <sub>3</sub>	I	H	H	H
	973	H	CF <sub>3</sub>	I	Cl	OH	H
25	974	H	CF <sub>3</sub>	I	Br	OH	H
	975	H	CF <sub>3</sub>	I	I	OH	H




- 712 -

5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	976	H	CF <sub>3</sub>	I	Br	H	H
	977	H	CF <sub>3</sub>	I	OH	H	H
	978	H	CF <sub>3</sub>	I	NO <sub>2</sub>	H	H
	979	H	CF <sub>3</sub>	I	I	H	H
	980	H	CF <sub>3</sub>	I	Cl	H	H
15	981	H	CF <sub>3</sub>	I	Cl	H	Cl
	982	H	CF <sub>3</sub>	I	OMe	H	H
	983	H	CF <sub>3</sub>	I	H	CF <sub>3</sub>	H
	984	H	CF <sub>3</sub>	I	H	OH	H
	985	H	CF <sub>3</sub>	I	H	OMe	H
20	986	H	CF <sub>3</sub>	I	H	Cl	H
	987	H	CF <sub>3</sub>	I	H	Cl	Cl
	988	H	CF <sub>3</sub>	I	H	Br	H
	989	H	CF <sub>3</sub>	I	CF <sub>3</sub>	H	H
	990	H	CF <sub>3</sub>	I	H	H	H
25	991	H	CF <sub>3</sub>	I	Cl	OH	H
	992	H	CF <sub>3</sub>	I	Br	OH	H
	993	H	CF <sub>3</sub>	I	I	OH	H

5



	<u>Example #</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
10	994	H	CF <sub>3</sub>	I	Br	H	H
	995	H	CF <sub>3</sub>	I	OH	H	H
	996	H	CF <sub>3</sub>	I	NO <sub>2</sub>	H	H
	997	H	CF <sub>3</sub>	I	I	H	H
	998	H	CF <sub>3</sub>	I	Cl	H	H
15	999	H	CF <sub>3</sub>	I	Cl	H	Cl
	1000	H	CF <sub>3</sub>	I	OMe	H	H
	1001	H	CF <sub>3</sub>	I	H	CF <sub>3</sub>	H
	1002	H	CF <sub>3</sub>	I	H	OH	H
	1003	H	CF <sub>3</sub>	I	H	OMe	H
20	1004	H	CF <sub>3</sub>	I	H	Cl	H
	1005	H	CF <sub>3</sub>	I	H	Cl	Cl
	1006	H	CF <sub>3</sub>	I	H	Br	H
	1007	H	CF <sub>3</sub>	I	CF <sub>3</sub>	H	H
	1008	H	CF <sub>3</sub>	I	H	H	H
25	1009	H	CF <sub>3</sub>	I	Cl	OH	H
	1010	H	CF <sub>3</sub>	I	Br	OH	H
	1011	H	CF <sub>3</sub>	I	I	OH	H

- 714 -

The activity of the compounds of the present invention was tested in the following assays. The results of testing in the assays are tabulated in Table 1.

5

#### VITRONECTIN ADHESION ASSAY

##### MATERIALS

Human vitronectin receptor ( $\alpha_v\beta_3$ ) was purified from human placenta as previously described [Pytela et al.,  
10 Methods in Enzymology, 144:475-489 (1987)]. Human vitronectin was purified from fresh frozen plasma as previously described [Yatohgo et al., Cell Structure and Function, 13:281-292 (1988)]. Biotinylated human vitronectin was prepared by coupling NHS-biotin from  
15 Pierce Chemical Company (Rockford, IL) to purified vitronectin as previously described [Charo et al., J. Biol. Chem., 266(3):1415-1421 (1991)]. Assay buffer, OPD substrate tablets, and RIA grade BSA were obtained from Sigma (St. Louis, MO). Anti-biotin  
20 antibody was obtained from Calbiochem (La Jolla, CA). Linbro microtiter plates were obtained from Flow Labs (McLean, VA). ADP reagent was obtained from Sigma (St. Louis, MO).

##### 25 METHODS

##### Solid Phase Receptor Assays

This assay was essentially the same as previously reported [Niiya et al., Blood, 70:475-483 (1987)]. The  
30 purified human vitronectin receptor ( $\alpha_v\beta_3$ ) was diluted from stock solutions to 1.0  $\mu\text{g/mL}$  in Tris-buffered saline containing 1.0 mM  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , and  $\text{Mn}^{++}$ , pH 7.4 (TBS $^{+++}$ ). The diluted receptor was immediately transferred to Linbro microtiter plates at 100  $\mu\text{L/well}$ .  
35 (100 ng receptor/well). The plates were sealed and incubated overnight at 4°C to allow the receptor to bind to the wells. All remaining steps were at room

- 715 -

temperature. The assay plates were emptied and 200  $\mu$ L of 1% RIA grade BSA in TBS<sup>+++</sup> (TBS<sup>+++</sup>/BSA) were added to block exposed plastic surfaces. Following a 2 hour incubation, the assay plates were washed with TBS<sup>+++</sup> using a 96 well plate washer. Logarithmic serial dilution of the test compound and controls were made starting at a stock concentration of 2 mM and using 2 nM biotinylated vitronectin in TBS<sup>+++</sup>/BSA as the diluent. This premixing of labeled ligand with test (or control) ligand, and subsequent transfer of 50  $\mu$ L aliquots to the assay plate was carried out with a CETUS Propette robot; the final concentration of the labeled ligand was 1 nM and the highest concentration of test compound was  $1.0 \times 10^{-4}$  M. The competition occurred for two hours after which all wells were washed with a plate washer as before. Affinity purified horseradish peroxidase labeled goat anti-biotin antibody was diluted 1:3000 in TBS<sup>+++</sup>/BSA and 125  $\mu$ L were added to each well. After 30 minutes, the plates were washed and incubated with OPD/H<sub>2</sub>O<sub>2</sub> substrate in 100 mM/L Citrate buffer, pH 5.0. The plate was read with a microtiter plate reader at a wavelength of 450 nm and when the maximum-binding control wells reached an absorbance of about 1.0, the final A<sub>450</sub> were recorded for analysis. The data were analyzed using a macro written for use with the EXCEL™ spreadsheet program. The mean, standard deviation, and %CV were determined for duplicate concentrations. The mean A<sub>450</sub> values were normalized to the mean of four maximum-binding controls (no competitor added) (B-MAX). The normalized values were subjected to a four parameter curve fit algorithm [Rodbard et al., Int. Atomic Energy Agency, Vienna, pp 469 (1977)], plotted on a semi-log scale, and the computed concentration corresponding to inhibition of 50% of the maximum binding of biotinylated vitronectin (IC<sub>50</sub>) and corresponding R<sup>2</sup> was reported for those compounds exhibiting greater than 50% inhibition at the

- 716 -

highest concentration tested; otherwise the  $IC_{50}$  is reported as being greater than the highest concentration tested.  $\beta$ -[[2-[[5-  
[(aminoiminomethyl)amino]-1-oxopentyl]amino]-1-oxoethyl]amino]-3-pyridinepropanoic acid [US 5 08/375,338, Example 1] which is a potent  $\alpha_v\beta_3$  antagonist ( $IC_{50}$  in the range 3-10 nM) was included on each plate as a positive control.

10 PURIFIED IIb/IIIa RECEPTOR ASSAY

MATERIALS

Human fibrinogen receptor ( $\alpha_{IIb}\beta_3$ ) was purified from outdated platelets. (Pytela, R., Pierschbacher, M.D.,  
15 Argraves, S., Suzuki, S., and Rouslahti, E. "Arginine-Glycine-Aspartic acid adhesion receptors", Methods in Enzymology 144(1987):475-489.) Human vitronectin was purified from fresh frozen plasma as described in  
Yatohgo, T., Izumi, M., Kashiwagi, H., and Hayashi, M.,  
20 "Novel purification of vitronectin from human plasma by heparin affinity chromatography," Cell Structure and Function 13(1988):281-292. Biotinylated human vitronectin was prepared by coupling NHS-biotin from  
Pierce Chemical Company (Rockford, IL) to purified  
25 vitronectin as previously described. (Charo, I.F., Nannizzi, L., Phillips, D.R., Hsu, M.A., Scarborough, R.M., "Inhibition of fibrinogen binding to GP IIb/IIIa by a GP IIIa peptide", J. Biol. Chem. 266(3)(1991):  
1415-1421.) Assay buffer, OPD substrate tablets, and  
30 RIA grade BSA were obtained from Sigma (St. Louis, MO). Anti-biotin antibody was obtained from Calbiochem (La Jolla, CA). Linbro microtiter plates were obtained from Flow Labs (McLean, VA). ADP reagent was obtained from Sigma (St. Louis, MO).

- 717 -

METHODSSolid Phase Receptor Assays

This assay is essentially the same reported in Niiya, K., Hodson, E., Bader, R., Byers-Ward, V. Koziol, J.A., Plow, E.F. and Ruggeri, Z.M., "Increased surface expression of the membrane glycoprotein IIb/IIIa complex induced by platelet activation: Relationships to the binding of fibrinogen and platelet aggregation", Blood 70(1987):475-483. The purified human fibrinogen receptor ( $\alpha_{IIb}\beta_3$ ) was diluted from stock solutions to 1.0  $\mu\text{g/mL}$  in Tris-buffered saline containing 1.0 mM  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , and  $\text{Mn}^{++}$ , pH 7.4 ( $\text{TBS}^{+++}$ ). The diluted receptor was immediately transferred to Linbro microtiter plates at 100  $\mu\text{L/well}$  (100 ng receptor/well). The plates were sealed and incubated overnight at 4°C to allow the receptor to bind to the wells. All remaining steps were at room temperature. The assay plates were emptied and 200  $\mu\text{L}$  of 1% RIA grade BSA in  $\text{TBS}^{+++}$  ( $\text{TBS}^{+++}/\text{BSA}$ ) were added to block exposed plastic surfaces. Following a 2 hour incubation, the assay plates were washed with  $\text{TBS}^{+++}$  using a 96 well plate washer. Logarithmic serial dilution of the test compound and controls were made starting at a stock concentration of 2 mM and using 2 nM biotinylated vitronectin in  $\text{TBS}^{+++}/\text{BSA}$  as the diluent. This premixing of labeled ligand with test (or control) ligand, and subsequent transfer of 50  $\mu\text{L}$  aliquots to the assay plate was carried out with a CETUS Propette robot; the final concentration of the labeled ligand was 1 nM and the highest concentration of test compound was  $1.0 \times 10^{-4}$  M. The competition occurred for two hours after which all wells were washed with a plate washer as before. Affinity purified horseradish peroxidase labeled goat anti-biotin antibody was diluted 1:3000 in  $\text{TBS}^{+++}/\text{BSA}$  and 125  $\mu\text{L}$  were added to each well. After 30 minutes, the

- 718 -

plates were washed and incubated with ODD/H<sub>2</sub>O<sub>2</sub> substrate in 100 mM/L citrat buffer, pH 5.0. The plate was read with a microtiter plate reader at a wavelength of 450 nm and when the maximum-binding control wells reached an absorbance of about 1.0, the final A<sub>450</sub> were recorded for analysis. The data were analyzed using a macro written for use with the EXCEL™ spreadsheet program. The mean, standard deviation, and %CV were determined for duplicate concentrations. The mean A<sub>450</sub> values were normalized to the mean of four maximum-binding controls (no competitor added) (B-MAX). The normalized values were subjected to a four parameter curve fit algorithm, [Robard et al., Int. Atomic Energy Agency, Vienna, pp 469 (1977)], plotted on a semi-log scale, and the computed concentration corresponding to inhibition of 50% of the maximum binding of biotinylated vitronectin (IC<sub>50</sub>) and corresponding R<sup>2</sup> was reported for those compounds exhibiting greater than 50% inhibition at the highest concentration tested; otherwise the IC<sub>50</sub> is reported as being greater than the highest concentration tested.  $\beta$ -[[2-[[5-[(aminoiminomethyl)amino]-1-oxopentyl]amino]-1-oxoethyl]amino]-3-pyridinepropanoic acid [USN 08/375,338, Example 1] which is a potent  $\alpha,\beta$  antagonist (IC<sub>50</sub> in the range 3-10 nM) was included on each plate as a positive control.

#### Human Platelet Rich Plasma Assays

Healthy aspirin free donors were selected from a pool of volunteers. The harvesting of platelet rich plasma and subsequent ADP induced platelet aggregation assays were performed as described in Zucker, M.B., "Platelet Aggregation Measured by the Photometric Method", Methods in Enzymology 169(1989):117-133. Standard venipuncture techniques using a butterfly allowed the withdrawal of 45 mL of whole blood into a 60 mL syringe containing 5 mL of 3.8% trisodium

- 719 -

citrate. Following thorough mixing in the syringe, the anti-coagulated whole blood was transferred to a 50 mL conical polyethylene tube. The blood was centrifuged at room temperature for 12 minutes at 200 xg to sediment non-platelet cells. Platelet rich plasma was removed to a polyethylene tube and stored at room temperature until used. Platelet poor plasma was obtained from a second centrifugation of the remaining blood at 2000 xg for 15 minutes. Platelet counts are typically 300,000 to 500,000 per microliter. Platelet rich plasma (0.45 mL) was aliquoted into siliconized cuvettes and stirred (1100 rpm) at 37°C for 1 minute prior to adding 50 uL of pre-diluted test compound. After 1 minute of mixing, aggregation was initiated by the addition of 50 uL of 200 uM ADP. Aggregation was recorded for 3 minutes in a Payton dual channel aggregometer (Payton Scientific, Buffalo, NY). The percent inhibition of maximal response (saline control) for a series of test compound dilutions was used to determine a dose response curve. All compounds were tested in duplicate and the concentration of half-maximal inhibition ( $IC_{50}$ ) was calculated graphically from the dose response curve for those compounds which exhibited 50% or greater inhibition at the highest concentration tested; otherwise, the  $IC_{50}$  is reported as being greater than the highest concentration tested.

#### M21 MELANOMA CELL ADHESION ASSAY

This assay involves an  $\alpha_v\beta_3$ -dependent adhesion of M21 human melanoma cells to human fibrinogen-coated plastic tissue culture dishes.

Fibrinogen was purified from human plasma. Fibronectin and plasminogen were eliminated from the preparation by passing the sample over gelatin-sepharose 4B and lysine-sepharose 4B resins,



- 720 -

- respectively. The fibrinogen is diluted to 10  $\mu\text{g/mL}$  in coating buffer (20 mM Tris-HCl, 150 mM NaCl, pH 7.4). 100  $\mu\text{L}$  of diluted fibrinogen is added to each well of a 96-well Immulon 2 microtiter plate (Dynatech;
- 5 Chantilly, Va) and allowed to coat overnight at 4°C. Plates are blocked with 1% BSA (Miles/Pentex; Kankakee, IL) in adhesion buffer (Hank's balanced salt solution without  $\text{Ca}^{++}$  or  $\text{Mg}^{++}$  [HBSS--], 50 mM Hepes, 1 mg/mL BSA, pH 7.4) for 1 hour at 37°C.
- 10 M21 human melanoma cells were provided by Dr. J. Smith, La Jolla Cancer Research Institute. M21 cells are harvested from tissue culture flasks by washing with HBSS-- and adding cell dissociation solution (Sigma) and incubating for 5 minutes at 37°C.
- 15 Harvested cells are washed 3 times with adhesion assay buffer containing 200  $\mu\text{M}$   $\text{Mn}^{++}$ . Cells are counted and suspended to a density of  $2 \times 10^6/\text{mL}$  in adhesion assay buffer containing 200  $\mu\text{M}$   $\text{Mn}^{++}$ . M21 cells are pre-incubated with antagonists of  $\alpha_v\beta_3$  for 30 minutes at
- 20 room temperature. Following the pre-incubation, the solutions containing a mixture of cells and antagonists are added to each well of the microtiter plate and allowed to bind for 30 minutes at 37°C.
- Following adhesion, plates are gently washed 3
- 25 times with 200  $\mu\text{L}$  of wash buffer (50 mM Tris-HCl, 150 mM NaCl, pH 7.4) using large bore pipet tips. Plates are briefly blotted dry and 100  $\mu\text{L}$  of cell lysis buffer (50 mM sodium acetate, pH 5.0, 0.5% Triton X-100, 0.3 mg/mL p-nitrophenyl phosphate [Sigma] is added to each
- 30 well. Plates are incubated for 60 minutes at 37°C and 50  $\mu\text{L}$  of 1N NaOH is added to stop the reaction. The absorbance of the wells at 412 nm is read using an automatic plate reader.

- 721 -

TABLE I

Example	AvB3 IC50 (nM)	IIb/IIIa IC50 (nM)	M21 Melanoma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
1	76.9	8350		> 200
2	0.54	51.2	0.25	200
3	498	72900	3050	
4	3.17	473	3.3	> 200
5	227	3150		
6	1.04	15.9		80
8	0.69	9.83	0.28	73.3
10	0.92	54.4	1.82	> 200
12	1.1	595	9.32	> 200
14	1.62	139	5.42	> 200
15	10.2	3830	202	> 200
17	2.66	137	3.64	> 200
19	303	72000		
21	2.44	1910		> 200
22	1.37	280		> 200
24	0.91	58.6	12.7	> 200
26	14.2	809		> 200
27	1.53	178		> 200
30	1.75	424	320	> 200
34	94.3	269		> 200
35	57.1	6.21		69.5
36 Step B	14.6	1580	143	> 200
37	0.88	13.9		> 20.0
39	12.2	1540		> 20.0
40	10.3	834		> 200
41	12.1	830		> 200
42	124	9800		
43	28.3	1640	188	> 200
44	0.33	998		> 20.0
45	0.69	39.5	2.54	167

- 722 -

Example	AvB3 IC50 (nM)	IIB/IIIA IC50 (nM)	M21 Melanoma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
46	5.34	1680	147	> 200
47	0.86	4270	1.18	> 200
51	9730	>100000		
52	3.62	139	11.7	> 200
53	54.6	930		> 200
54	10.7	175		> 200
55	4.77	117		> 200
56	3.12	65.3	6.87	> 200
57	1340	15300		
58	162	5740		
59	2.35	172	24.3	> 200
60(B)	1.21	72.7		> 200
60(C)	0.73	16.4	0.74	> 200
61	1.76	192	228	> 200
62	1.42	28.4		> 200
65	9.7	170	13.8	> 200
66	1.44	73.7	2.51	> 100
67	2.05	92.3	4.08	> 200
68	5.48	125		> 200
69	0.92	33.6	0.95	> 200
70	63	3240	924	> 200
71	20.4	202	1040	> 200
72	1.21	152		> 200
80	9.49	4.35		30
82	334	353		
83	3.39	97.7	11	> 200
84	2800	246		
85	6.65	8.07		
86	8.79	246		> 200
87	6.35	732		> 200
88	8.44	945	52.3	> 200

- 723 -

Example	AvB3 IC50 (nM)	IIb/IIIa IC50 (nM)	M21 Melanoma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
89	1240	9830		
94	1.16	101	1	> 200
95	1.43	25.4		> 200
96	1810	5400		
97	26.9	1170	163	
98	146	500		
99	0.38	1.89	0.49	57.5
100	8560	>100000		
101	1680	65700		
103	16.6	19100		> 20.0
106	0.79	3140	0.81	> 200
107	6400	18700		
108	25.2	4870		> 200
109	575	>100000		
110	4.5	1860	177	> 200
112	284	6340		
113	276	100000		
114	3.26	2940	200	> 200
116	15500	>100000		
117	60.1	20100		> 200
119	3.61	11100	90.4	> 20.0
121	2840	>100000		
122	0.79	420		> 20.0
123	11800	85500		
124	22	317		> 20.0
126	2.48	2010		> 200
127	0.51	461		> 200
129	68.9	9460		> 200
130	47	2690		> 200
131	3.82	1760		> 20.0
135	50700	>100000		

- 724 -

Example	AvB3 IC50 (nM)	IIb/IIIa IC50 (nM)	M21 Melanoma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
136	54.4	14200		> 20.0
137	16.2	6500		> 200
138	36.9	5820		> 200
139	23.8	16100		> 200
140	4590	>100000		
141	3.09	125		> 200
143	6700	>100000		
144	55.3	5830		> 200
145	2720	>100000		
146	14.3	879		> 200
150	5.74	631		> 200
155	5.05	81.1		> 200
158	10.1	547		
160	25.6	10400		
162	4.62	1340		>200
166	13000	45900		
168	2.29	269		
171	0.35	83.2		
173	0.5	17.4		
175	2.12	205		
177	0.58	137		>20.0
179	2.72	927		
181	132	22800		
183	1.58	258		
185	1.47	166		
187	1.31	264		
189	4.03	1980		
191	0.49	70.3		>20.0
193	2.56	209		>20.0
195	1.09	98		
198	114	37800		
200	0.48	1100		>200

- 725 -

Example	AvB3 IC50 (nM)	I Ib/IIIa IC50 (nM)	M21 Melanoma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
201	58.1	10800		
203	3.56	650		
205	1.68	1240		
206	78.5	22000		
207	0.9	148		
208	1.15	277		
209	0.83	140		
210	2.62	343		
211	0.47	607		
212	1.93	306		
213	2.93	334		
214	2.35	454		
215	0.41	656		
216	1	326		
217	74.8	78900		
219	2.29	253		
221	70.5	23.7		>200
222	2.02	112		>200
223	4.36	293		>200
224	0.71	25.9		
225	2.76	471		>20.0
226	7.07	2910		>200
227	14.1	2640		>200
228	3.36	583		>200
229	39.1	10600		
231	2.99	424		
232	19.1	12100		>200
233	3.31	647		>200
234	89.3	830		
235	0.54	29.9		
236	0.53	1250		
237	0.57	1950		
238	0.92	646		

- 726 -

Example	AvB3 IC50 (nM)	IIb/IIIa IC50 (nM)	M21 Melan ma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
239	0.83	673		
240	49400	76400		
241	557	17200		
242	2.28	533		
243	0.35	23.6		
244	17.6	4560		
245	0.96	134		
246	7.24	802		
247	1.24	417		
248	12300	21000		
249	5.31	244		
251(B)	3.49	280		
251(C)	0.76	124		
252	1.52	213		
253	0.84	109		
254	16.5	6910		
255	28.4	6050		
256	0.58	22		
257	49.2	4660		
259	0.81	86.7		
260	0.74	65.3		
261	6.47	4710		
262	1.24	172		
263	4.19	2760		
264	2.18	574		
265	6.19	706		
266	0.77	1810		
267	131	43900		
268	0.67	7430		
269	209	25400		
270	5.51	9160		
271	29.9	4610		
272	893	8210		

- 727 -


Example	AvB3 IC50 (nM)	IIb/IIIa IC50 (nM)	M21 M lan ma Cells IC 50 (nM)	Human PRP ( $\mu$ M)
273	12.9	4160		
274	31.1	21200		
275	6.98	1200		
276	1.25	111		
277	1.41	198		
278	0.45	150		
279	7.12	637		
281	4.16	11500		
282	864	9770		
284	195	18400		
285	229	3170		
286	413	8090		
287	49.7	41.1		
288	8.62	1060		
289	0.9	621		
290	1.62	1020		
291	1.24	37.4		
292	3.55	337		
294	173	1990		
295	144	4560		
296	404	9450		
297	89.8	3920		
298	252	5560		
299	109	927		
362	0.84	7260		
363	2.12	509		
364	3.58	223		
365	16.9	8470		
366	0.44	91.3		
367	0.35	1540		



- 728 -


What is claimed is:

1. A compound of the formula



or a pharmaceutically acceptable salt thereof,  
wherein

A is



wherein Y<sup>1</sup> is selected from the group consisting of  
N-R<sup>2</sup>, O, and S;

R<sup>2</sup> is selected from the group consisting of H;  
alkyl; aryl; hydroxy; alkoxy; cyano; nitro; amino;  
alkenyl; alkynyl; amido; alkylcarbonyl;  
arylcarbonyl; alkoxycarbonyl; aryloxycarbonyl;  
haloalkylcarbonyl; haloalkoxycarbonyl;  
alkylthiocarbonyl; arylthiocarbonyl;  
acyloxymethoxycarbonyl; alkyl optionally  
substituted with one or more substituent selected  
from lower alkyl, halogen, hydroxyl, haloalkyl,  
cyano, nitro, carboxyl, amino, alkoxy, aryl or  
aryl optionally substituted with one or more  
halogen, haloalkyl, lower alkyl, alkoxy, cyano,  
alkylsulfonyl, alkylthio, nitro, carbonyl, amino,  
hydroxyl, sulfonic acid, sulfonamide, aryl, fused  
aryl, monocyclic heterocycles, or fused monocyclic

- 729 -

heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, hydroxy, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, cyano, nitro, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, carboxyl derivatives, amino, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycle; monocyclic heterocycles; and monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, aryl or fused aryl; or

$R^2$  taken together with  $R^7$  forms a 4-12 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, alkoxy, halo, phenyl, amino, carboxyl or carboxyl ester, and fused phenyl;

or  $R^2$  taken together with  $R^7$  forms a 5 membered heteroaromatic ring optionally substituted with one or more substituent selected from lower alkyl, phenyl and hydroxy;

or  $R^2$  taken together with  $R^7$  forms a 5 membered heteroaromatic ring fused with a phenyl group;


$R^7$  (when not taken together with  $R^2$ ) and  $R^8$  are independently selected from the group consisting of H; alkyl; alkenyl; alkynyl; aralkyl; amino; alkylamino; hydroxy; alkoxy; arylamino; amido, alkylcarbonyl, arylcarbonyl; alkoxycarbonyl; aryloxy; haloalkylcarbonyl; aryloxy; haloalkoxycarbonyl; alkylthiocarbonyl;

- 730 -

arylthiocarbonyl; acyloxymethoxycarbonyl; cycloalkyl; bicycloalkyl; aryl; acyl; benzoyl; alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxy, haloalkyl, cyano, nitro, carboxyl derivatives, amino, alkoxy, thio, alkylthio, sulfonyl, aryl, aralkyl, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethyl, sulfonyl, alkylsulfonyl, haloalkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, fused monocyclic heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles; monocyclic heterocycles; monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, aryloxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, aryl, fused aryl; monocyclic and bicyclic heterocyclicalkyls;  $-SO_2R^{10}$  wherein  $R^{10}$  is selected from the group consisting of alkyl, aryl and monocyclic heterocycles, all optionally substituted with one or more substituent selected from the group consisting of halogen, haloalkyl,

- 731 -


alkyl, alkoxy, cyano, nitro, amino, acylamino, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, trifluoromethylthio, trifluoroalkoxy, trifluoromethylsulfonyl, aryl, aryloxy, thio, alkylthio, and monocyclic heterocycles; and



or NR<sup>7</sup> and R<sup>8</sup> taken together form a 4-12 membered mononitrogen containing monocyclic or bicyclic ring optionally substituted with one or more substituent selected from lower alkyl, carboxyl derivatives, aryl or hydroxy and wherein said ring optionally contains a heteroatom selected from the group consisting of O, N and S;

R<sup>5</sup> is selected from the group consisting of H, alkyl, alkenyl, alkynyl, benzyl, and phenethyl;

or



wherein Y<sup>2</sup> is selected from the group consisting of alkyl; cycloalkyl; bicycloalkyl; aryl; monocyclic heterocycles; alkyl optionally substituted with aryl which can also be optionally substituted with one or more substituent selected from halo, haloalkyl, alkyl, nitro, hydroxy, alkoxy, aryloxy, aryl, or fused aryl; aryl optionally substituted


- 732 -

with one or more substituent selected from halo, haloalkyl, hydroxy, alkoxy, aryloxy, aryl, fused aryl, nitro, methylenedioxy, ethylenedioxy, or alkyl; alkynyl; alkenyl;  $-S-R^9$  and  $-O-R^9$  wherein  $R^9$  is selected from the group consisting of H; alkyl; aralkyl; aryl; alkenyl; and alkynyl; or  $R^9$  taken together with  $R^7$  forms a 4-12 membered mononitrogen and monosulfur or monooxygen containing heterocyclic ring optionally substituted with lower alkyl, hydroxy, keto, phenyl, carboxyl or carboxyl ester, and fused phenyl; or  $R^9$  taken together with  $R^7$  is thiazole; oxazole; benzoxazole; or benzothiazole; and

$R^5$  and  $R^7$  are as defined above;

or  $Y^2$  (when  $Y^2$  is carbon) taken together with  $R^7$  forms a 4-12 membered mononitrogen containing ring optionally substituted with alkyl, aryl or hydroxy;

or A is




where  $R^2$  and  $R^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, phenyl, or carboxyl derivatives; and  $R^8$  is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, or acyloxymethoxycarbonyl; and

- 733 -

$R^5$  is defined as above

or A is



where  $R^2$  and  $R^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with hydroxy, keto, phenyl, or alkyl; and

$R^8$  are both selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl and acyloxymethoxycarbonyl;

$Z^1$  is one or more substituent selected from the group consisting of H; alkyl; hydroxy; alkoxy; aryloxy; halogen; haloalkyl; haloalkoxy; nitro; amino; alkylamino; acylamino; dialkylamino; cyano; alkylthio; alkylsulfonyl; carboxyl derivatives; trihaloacetamide; acetamide; aryl; fused aryl; cycloalkyl; thio; monocyclic heterocycles; fused monocyclic heterocycles; and A, wherein A is defined above;

V is selected from the group consisting of  $-N-(R^6)-$  wherein  $R^6$  is selected from the group consisting of H; lower alkyl; cycloalkyl; aralkyl; aryl; and monocyclic heterocycles; or  $R^6$  taken together with Y, forms a 4-12 membered mononitrogen containing ring;

- 734 -

Y, Y<sup>3</sup>, Z and Z<sup>3</sup> are independently selected from the group consisting of hydrogen; alkyl; aryl; and cycloalkyl; or Y and Z taken together form a cycloalkyl; or Y<sup>3</sup> and Z<sup>3</sup> taken together form a cycloalkyl;

n is an integer 1, 2, or 3;

t is an integer 0, 1, or 2;

p is an integer 0, 1, 2, or 3;

R is X-R<sup>3</sup> wherein X is selected from the group consisting of O, S and NR<sup>4</sup>, wherein R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydrogen; alkyl; alkenyl; alkynyl; haloalkyl; aryl; arylalkyl; sugars; steroids; polyalkylethers; alkylamido; alkyl N,N-dialkylamido; pivaloyloxymethyl; and in the case of the free acid, all pharmaceutically acceptable salts thereof;

R<sup>1</sup> is selected from the group consisting of hydrogen; alkyl; alkenyl; alkynyl; aryl; carboxyl derivatives; haloalkyl; monocyclic heterocycles; monocyclic heterocycles optionally substituted with alkyl, halogen, haloalkyl, cyano, hydroxy, aryl, fused aryl, nitro, alkoxy, aryloxy, alkylsulfonyl, arylsulfonyl, sulfonamide, thio, alkylthio, carboxyl derivatives, amino, amido;


alkyl optionally substituted with one or more of halo, haloalkyl, hydroxy, alkoxy, aryloxy, thio, alkylthio, alkynyl, alkenyl, alkyl, arylthio, alkylsulfoxide, alkylsulfonyl, arylsulfoxide, arylsulfonyl, cyano, nitro, amino, alkylamino, dialkylamino, alkylsulfonamide, arylsulfonamide,

- 735 -

acylamide, carboxyl derivatives, sulfonamide, sulfonic acid, phosphonic acid derivatives, phosphinic acid derivatives, aryl, arylthio, arylsulfoxide, or arylsulfone all optionally substituted on the aryl ring with halo, alkyl, haloalkyl, cyano, nitro, hydroxy, carboxyl derivatives, alkoxy, aryloxy, amino, alkylamino, dialkylamino, amido, aryl, fused aryl, monocyclic heterocycles; and fused monocyclic heterocycles, monocyclic heterocyclicthio, monocyclic heterocyclicsulfoxide, and monocyclic heterocyclic sulfone, which can be optionally substituted with halo, haloalkyl, nitro, hydroxy, alkoxy, fused aryl, or alkyl;

alkylcarbonyl, haloalkylcarbonyl, and arylcarbonyl;

aryl optionally substituted in one or more positions with halo, haloalkyl, alkyl, alkoxy, aryloxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, acyloxy, carboxyl derivatives, carboxyalkoxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycles; and



wherein R<sup>7</sup> and R<sup>8</sup> are as defined above

and provided that taken together with the nitrogen, R<sup>7</sup> and R<sup>8</sup> comprise an amino acid;

and




- 736 -

R<sup>11</sup> is selected from the group consisting of H, alkyl, aralkyl, alkenyl, alkynyl, haloalkyl or haloalkynyl or R<sup>11</sup> taken together with Y forms a 4-12 membered mononitrogen containing ring.

2. A compound according to the formula of Claim 1 wherein

A is



wherein Y<sup>1</sup> is selected from the group consisting of N-R<sup>2</sup>, O, and S;

R<sup>2</sup> is selected from the group consisting of H, cyano, alkyl, aryl, substituted alkyl, hydroxy, alkoxy, alkylcarbonyl, amido, nitro, amino and monocyclic heterocycles, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl; or R<sup>2</sup> taken together with R<sup>7</sup> forms a 4-12 membered ring;

R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup> are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, bicycloalkyl substituted alkyl, arylalkyl, aryloxy, hydroxy, alkoxy, amino, alkylamino, arylamino, amido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl, substituted phenyl,

- 737 -

arylacyl, monocyclic and bicyclic heterocycles, monocyclic and bicyclic heterocyclicalkyl and -SO<sub>2</sub>R<sup>10</sup> wherein R<sup>10</sup> is selected from the group consisting of alkyl, amino and aryl which are all optionally substituted with one or more substituent selected from the group consisting of acylamino, amino, carbonyl, cyano, nitro, alkoxy, halo, alkyl, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, aryloxy, thio, trifluoromethylthio, trifluoroalkoxy, and trifluoromethylsulfonyl or -NR<sup>7</sup> and R<sup>8</sup> taken together form a 4-12 membered ring wherein said ring optionally contains a heteroatom selected from the group consisting of O, N, and S wherein said ring is optionally substituted;

3. A compound according to Claim 2 wherein

V is -N(R<sup>6</sup>)- wherein R<sup>6</sup> is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0 or 1;

p is 0, 1 or 2; and

R is O-R<sup>3</sup>.

4. A compound according to Claim 3 selected from the group consisting of

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

- 738 -

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-  
propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]benzenepropanoate;

(±)β-[[2-[[[3-  
[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
naphthalen-1-yl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)-amino]-  
naphthalen-1-yl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

(±)ethyl β-[[1-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]-1-cyclopropyl]-  
carbonyl]amino]pyridine-3-propanoate;

(±)β-[[1-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]cycloprop-1-yl]carbonyl]-  
amino]pyridine-3-propanoic acid;

- 739 -

(±) ethyl  $\beta$ -[[2-[[[3-[[imino[(phenylmethyl)amino]-methyl]amin ]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[[imino[(phenylmethyl)amino]methyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

3S-[[2-[[[3-(aminocarbonylamino)phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-(aminocarbonylamino)phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoate;

$\beta$ S-[[2-[[[3-[(aminoiminomethyl)amino]-2,5,6-trifluorophenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

ethyl (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propanoic acid;

- 740 -

ethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propanoate;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-naphthalene-1-carboxylic acid;

methyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-naphthalene-1-carboxylate;

( $\pm$ ) 3-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]-2-oxopyrrolidine-1-propanoic acid;

3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoate;

- 741 -

3S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-p ntynoic acid;

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]furan-2-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
furan-3-propanoic acid;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]pentanedioic acid;

bismethyl ester 3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]-amino]acetyl]amino]pentanedioate;

( $\pm$ ) hydrogen methyl 3-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pentanedioate;

- 742 -

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
furan-2-propanoic acid;

ethyl (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]furan-2-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
naphthalene-2-propanoic acid;

(±) methyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
thiophene-3-propanoate;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-pentynoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
thiophene-3-propanoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
carboxybutyl]thio]benzoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
carboxybutyl]sulfonyl]benzoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
thiophene-2-propanoic acid;

- 743 -

(±) methyl 2-[[3-[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-carboxybutyl]thio]benzoate;

(±) methyl 3-[[2-[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)thio]pentanoate;

(±) methyl 3-[[2-[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-[[4-methylphenyl)sulfonyl]amino]butanoate;

3-[[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-[(4-methylphenyl)sulfonyl]amino]butanoic acid;

(±) 3-[[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)thio]pentanoic acid;

(±) 3-[[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)sulfonyl]pentanoic acid;

3S-[[2-[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-(phenylthio)butanoic acid;

3R-[[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;



- 744 -

2-[[2S-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-2-  
(carboxymethyl)ethyl]sulfonyl]benzoic acid;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-pentynoate;

2-[[2S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-2-  
(carboxymethyl)ethyl]thio]benzoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]methylamino]acetyl]-  
amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]methylamino]acetyl]amino]pyridine-  
3-propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]-1-oxopropyl]-  
amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]-1-oxopropyl]-  
amino]pyridine-3-propanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-4-  
methylphenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

- 745 -

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-methyl]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-hydroxybutanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-2-hydroxy-5-methylbenzenepropanoic acid;

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-[(2-hydroxyethyl)amino]-4-oxobutanoic acid;

2S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-carboxypropyl 2-aminobenzoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-1,4-benzodioxin-6-propanoic acid;

N-[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]-β-alanine, ethyl ester;

N-[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]-β-alanine;

- 746 -

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-quinoline-3-propanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-quinoline-3-propanoic acid;

N-[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]-β-alanine, ethyl ester;

3-[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]amino]propanoic acid;

(±)ethyl β-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±) β-[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

ethyl β-[[2-[[[3-[[[(phenylamino)carbonyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

- 747 -

$\beta$ -[[2-[[[3-[(phenylamino)carbonyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]-amino]carbonyl]amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminothioxomethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

- 748 -

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-3-[[[(phenylamino)carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-3-[[[(phenylamino)carbonyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoic acid;

$\beta$ -[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]pyridine-  
3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(3-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(2-carboxyethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amin]-  
pyridine-3-propanoic acid;

- 749 -

$\beta$ -[[2-[[[3-[[[(2-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(1-naphthalenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
chlorophenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
chlorophenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-chlorophenyl]-  
carbonyl]amino]acetyl]amino]benzenepropanoic acid;

ethyl  $\beta$ -[2-[[[3-[[amino(aminocarbonyl)-  
imino]methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(aminocarbonyl)imino]-  
methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzene-  
propanoic acid;

[(dimethylamino)carbonyl]methyl  $\beta$ -  
[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoate;

1,1-dimethylethyl 3,5-dichloro- $\beta$ -[[2-[[[3[[[(ethoxy-  
carbonyl)amino]amino]amino]amino]amino]amino]amino]amino]-  
phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

- 750 -

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(ethoxycarbonyl)-amino]([thoxycarbonyl]imino)methyl]-amino]-phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoate;

3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]-methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)-amino]-4-chlorophenyl]carbonyl]amino]-acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoic acid;

- 751 -

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-bromobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-bromobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-bromobenzenepropanoate;



- 752 -

- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-bromobenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethylbenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethylbenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethoxybenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethoxybenzenepropanoic acid;
- (±) (2,2-dimethyl-1-oxopropoxy)methyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[[[(aminocarbonyl)imino)-methylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,4-dibromobenzenepropanoic acid;

- 753 -

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-fluoro-5-(trifluoromethyl)benzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-fluorobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-methylbenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-chlorobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-chlorobenzenepropanoic acid;

(±) [2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]-ethyl]  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-iodobenzenepropanoic acid;

- 754 -

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2-hydroxy-4-methoxybenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-hydroxy-4-methoxybenzofuran-6-propanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-2-hydroxybenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2-hydroxy-5-nitrobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoic acid;

- 755 -

thyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-cyclohexanepropanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-cyclohexanepropanoate;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichloro-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-chloro-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-chloro-2-hydroxybenzenepropanoic acid;

(±) 5-amino-β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromopyridine-3-propanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromopyridine-3-propanoate;

- 756 -

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(ethoxycarbonyl)amino]-thioxomethyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoic acid;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(ethoxycarbonyl)amino]-iminomethyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino][1,1'-biphenyl]-3-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino][1,1'-biphenyl]-3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-amino]acetyl]amino][pyrimidine-5-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-[pyrimidine-5-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-amino]acetyl]amino]-3-methylthiophene-2-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3-methylthiophene-2-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-(methylthio)-benzenepropanoic acid;

- 757 -

1,1-dimethylethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3-(methylthio)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-6-methylpyridine-2-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-(methylsulfonyl)-benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-diethoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-diethoxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromothiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromothiophene-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-chlorothiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-chlorothiophene-2-propanoate;

- 758 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1H-pyrazole-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1H-pyrazole-  
3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-methylthiophene-  
2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-methylthiophene-  
2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2,3,5-trichloro-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2,3,5-trichloro-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoic acid;

- 759 -

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-6-chloro-  
1,3-benzodioxole-5-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-6-chloro-  
1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzofuran-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzofuran-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3-  
(carboxymethoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3-  
(carboxymethoxy)benzenepropanoate;



- 760 -

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4,4,4-  
trifluorobutanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4,4,4-  
trifluorobutanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-bromo-4,5-  
dimethoxybenzenepropanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-bromo-4,5-  
dimethoxybenzenepropanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
methylpentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
methylpentanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pentanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
5-bromo-3-chloro-2-hydroxybenzene-  
propanoic acid;

- 761 -

$\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

- 762 -

ethyl 3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]pyridine-3-  
propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

3-[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]butanoic acid;

ethyl 3-[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]butanoate;

- 763 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-bis(1-methoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-bis(1-methoxy)benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dibromo-4-hydroxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dibromo-4-hydroxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-4-hydroxybenzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-4-hydroxybenzenepropanoate;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-hydroxyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-hydroxyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(phenoxyamino)carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

- 764 -

ethyl  $\beta$ -[[2-[[[3-[[ (phenoxyamino) carbonyl] amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[ (phenylamino) amino] carbonyl]-  
amino] phenyl] carbonyl] amino] acetyl] amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[ (phenylamino) amino] carbonyl]-  
amino] phenyl] carbonyl] amino] acetyl] amino]-  
pyridine-3-propanoate;

3-[[2-[[[3-[(aminoiminomethyl) amino] phenyl]-  
carbonyl] amino] acetyl] amino]-5-[(3,5-  
dichlorophenyl) amino]-5-oxopentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl) amino] phenyl]-  
carbonyl] amino] acetyl] amino]-5-[(3,5-  
dichlorophenyl) amino]-5-oxopentanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl) amino]-5-carboxy-  
phenyl] carbonyl] amino] acetyl] amino] pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl) amino]-5-carboxy-  
phenyl] carbonyl] amino] acetyl] amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl) amino]-5-carboxy-  
phenyl] carbonyl] amino] acetyl] amino]-3,5-  
dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl) amino]-5-carboxy-  
phenyl] carbonyl] amino] acetyl] amino]-3,5-  
dichlorobenzenepropanoate;

- 765 -

$\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dichloro-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-dichloro-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-  
(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-  
(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-(acetylamino)-5-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-(acetylamino)-5-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
3,5-dichlorobenzenepropanoate;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-  
[[[(methylamino)(methylimino)methyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-  
[[[(ethylamino)(methylimino)methyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

- 766 -

(±) 3,5-dichloro-β-[[2-[[[3-[[[(1-methylethyl)amino](methylinino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) β-[[2-[[[3-[(ethylamino)-(methylinino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]-4-fluorobenzene-propanoic acid;

(±) 4-fluoro-β-[[2-[[[3-[[[(4-pyridinylmethyl)amino](methylinino)methyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-fluorobenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-1H-imidazole-2-propanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2,3,4,6-tetrafluorobenzenepropanoic acid;

β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromothiophene-2-propanoic acid;

β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-2-mercaptobenzenepropanoic acid;

and

- 767 -

(±)  $\beta$ -[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-5-chloro-2-mercaptobenzenepropanoic acid.

5. A compound according to Claim 3 wherein  $Y^1$  is  $N-R^2$  and  $R^2$  is cyano.
6. A compound according to Claim 5 wherein the compound is selected from the group consisting of

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)-phenylmethylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)-methylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)-(amino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)(ethylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)(phenylmethyl)amino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(cyanoimino)(methylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[amino(cyanoimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;



- 768 -

$\beta$ -[[2-[[[3-[(cyanoimino)(ethylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

ethyl 3S-[[2-[[[3-[(cyanoimino)-  
(methylamino)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(cyanoimino)(methylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoic acid;

ethyl  $\beta$ -[[2-[[[3-[(cyanoimino)[2-pyridinylmethyl]-  
amino]methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)[2-pyridinylmethyl]-  
amino]methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(cyanoimino)[3-  
pyridinylmethyl]amino]methyl]amino]phenyl]-  
carbonyl]amino]acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)[3-pyridinylmethyl]-  
amino]methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[[(4-(aminosulfonyl)-  
phenylmethyl]amino)(cyanoimino)methyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoate;

3S-[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl]-  
amino)(cyanoimino)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-4-pentynoic acid;

- 769 -

ethyl  $\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

( $\pm$ ) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)-(methylamino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate;


( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)(methylamino)-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

and

ethyl 3-[[2-[[[3-[[amino(cyanoimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate.

7. A compound according to Claim 2 wherein

A is



wherein  $Y^1$  is  $N-R^2$ ;  $R^2$  taken together with  $R^7$  forms a 4-12 membered ring; and  $R^5$  and  $R^8$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, bicycloalkyl substituted alkyl, arylalkyl, aryloxy, hydroxy, alkoxy, amino, alkylamino, arylamino, amido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, haloalkylcarbonyl, haloalkoxycarbonyl,

- 770 -

alkylthiocarbonyl, arylthiocarbonyl, acyloxymethyl carbonyl, substituted phenyl, arylacyl, monocyclic and bicyclic heterocycles, monocyclic and bicyclic heterocyclicalkyl and -SO<sub>2</sub>R<sup>10</sup> wherein R<sup>10</sup> is selected from the group consisting of alkyl, amino and aryl which are all optionally substituted with one or more substituent selected from the group consisting of acylamino, amino, carbonyl cyano, nitro, alkoxy, halo, alkyl, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, aryloxy, thio, trifluoromethylthio, trifluoroalkoxy, and trifluoromethylsulfonyl or -NR<sup>7</sup> and R<sup>8</sup> taken together form a 4-12 membered ring wherein said ring optionally contains a heteroatom selected from the group consisting of O, N, and S wherein said ring is optionally substituted.

8. A compound according to Claim 7 wherein

V is -N(R<sup>6</sup>)- wherein R<sup>6</sup> is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0; and

p is 1.

- 771 -

9. A compound according to Claim 8 selected from the group consisting of

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

- 772 -

(±) 3,5-dibromo-β-[[2-[[[3-[(1,4,5,6-tetrahydr pyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

(±) 3-bromo-5-chloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) [2-[2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) [2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3-bromo-5-chloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

ethyl (±) 3-bromo-5-chloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

- 773 -

(±) 3,5-dichloro-β-[[2-[[[3-[(4,5-dihydro-1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]-amino]acetyl]amino]-2-hydroxybenzenepropanoic acid;

(±) 3-bromo-5-chloro-β-[[2-[[[3-[(4,5-dihydro-1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]-2-hydroxybenzenepropanoic acid;

β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

(±) ethyl 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

ethyl (±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;


- 774 -

(±) 3-bromo-5-dichloro-2-hydroxy-  
 $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-  
 dimethylpyrimidin-2-yl)amino]phenyl]-  
 carbonyl]amino]acetyl]amino]benzene-  
 propanoic acid;

and

ethyl(±) 3-bromo-5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-  
 [(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-  
 yl)amino]phenyl]carbonyl]amino]acetyl]-  
 amino]benzenepropanoate.

10. A compound according to the formula of Claim 1  
 wherein



wherein Y<sup>2</sup> is selected from the group consisting of  
 lower alkyl, substituted alkyl, phenyl,  
 substituted phenyl, cycloalkyl, monocyclic  
 heterocycles, -S-R<sup>9</sup> and -O-R<sup>9</sup> wherein R<sup>9</sup> is selected  
 from the group consisting of H, alkyl, substituted  
 alkyl, phenyl, substituted phenyl and monocyclic  
 heterocycles or R<sup>9</sup> taken together with R<sup>7</sup> forms a  
 4-12 membered ring; or

Y<sup>2</sup> taken together with R<sup>7</sup> forms a 4-12 membered  
 ring which is optionally substituted.

- 775 -

11. A compound according to Claim 10 wherein

$Y^2$  taken together with  $R^7$  forms a 4-12 membered ring which is optionally substituted.

12. A compound according to Claim 11 wherein

$V$  is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H and lower alkyl;

$n$  is 1;

$t$  is 0 or 1; and

$p$  is 1.

13. A compound according to Claim 12 wherein the compound is selected from the group consisting of

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoic acid;



- 776 -

- (±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;
- $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoate;
- (±)  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoic acid;
- (±)  $\beta$ -[[2-[[[3,5-bis[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;
- $\beta$ S-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoic acid;

- 777 -

(±) ethyl β-[[2-[[[3-[(3,4-dihydro-2H-pyrr 1-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;

(±) β-[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro-β-[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro-β-[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

(±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

and

ethyl(±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate.

14. A compound according to Claim 10 wherein

Y<sup>2</sup> is selected from the group consisting of lower alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl and monocyclic heterocycles.

- 778 -

15. A compound according to Claim 14 wherein

V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0; and

p is 1.

16. A compound according to Claim 15 wherein the compound is selected from the group consisting of

(±) ethyl  $\beta$ -[[2-[[[3-[(iminophenylmethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ S-[[2-[[[3-[[imino(1-pyrrolidinyl)methyl]-amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

(±)  $\beta$ -[[2-[[[3-[(iminophenylmethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

3,5-dichloro- $\beta$ -[[2-[[[3-[[imino(1-piperidinyl)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

and

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[imino(1-piperidinyl)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate.

- 779 -

17. A compound according to Claim 10 wherein  $Y^2$  is  $-S-R^9$  or  $-O-R^9$  wherein  $R^9$  is selected from the group consisting of H, alkyl, substituted alkyl, phenyl, substituted phenyl and monocyclic heterocycles or  $R^9$  taken together with  $R^7$  forms a 4-12 membered ring.

18. A compound according to Claim 17 wherein

V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H, lower alkyl, substituted alkyl, cycloalkyl, aryl, substituted aryl, monocyclic heterocycles and benzyl;

n is 0;

t is 0; and

p is 1 or 2.

19. A compound according to Claim 18 wherein the compound is selected from the group consisting of

ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydrothiazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(4,5-dihydrothiazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[(benzoxazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(benzoxazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate.




- 780 -

$\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;





and

ethyl  $\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate.






20. A compound according to Claim 1 of the formula








- 781 -



- 782 -







- 783 -













- 784 -







- 785 -








- 786 -







- 787 -







- 788 -







- 789 -



- 790 -






- 791 -












- 792 -






- 793 -






- 794 -





- 795 -






- 796 -






- 797 -



- 798 -






- 799 -











- 800 -







- 801 -





- 802 -




- 803 -



- 804 -




21. A pharmaceutical composition comprising a therapeutically effective amount of a compound of the formula



or a pharmaceutically acceptable salt thereof,  
wherein

A is



wherein Y<sup>1</sup> is selected from the group consisting of N-R<sup>2</sup>, O, and S;

- 805 -

$R^2$  is selected from the group consisting of H; alkyl; aryl; hydroxy; alkoxy; cyano; nitro; amino; alkenyl; alkynyl; amido; alkylcarbonyl; arylcarbonyl; alkoxycarbonyl; aryloxy carbonyl; haloalkylcarbonyl; haloalkoxycarbonyl; alkylthiocarbonyl; arylthiocarbonyl; acyloxymethoxycarbonyl; alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxyl, haloalkyl, cyano, nitro, carboxyl, amino, alkoxy, aryl or aryl optionally substituted with one or more halogen, haloalkyl, lower alkyl, alkoxy, cyano, alkylsulfonyl, alkylthio, nitro, carboxyl, amino, hydroxyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, hydroxy, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, cyano, nitro, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, carboxyl derivatives, amino, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycle; monocyclic heterocycles; and monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, aryl or fused aryl; or

$R^2$  taken together with  $R^1$  forms a 4-12 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, alkoxy, halo, phenyl, amino, carboxyl or carboxyl ester, and fused phenyl;

- 806 -


or  $R^2$  taken together with  $R^7$  forms a 5 membered heteroaromatic ring optionally substituted with one or more substituent selected from lower alkyl, phenyl and hydroxy;

or  $R^2$  taken together with  $R^7$  forms a 5 membered heteroaromatic ring fused with a phenyl group;

$R^7$  (when not taken together with  $R^2$ ) and  $R^8$  are independently selected from the group consisting of H; alkyl; alkenyl; alkynyl; aralkyl; amino; alkylamino; hydroxy; alkoxy; arylamino; amido, alkylcarbonyl, arylcarbonyl; alkoxycarbonyl; aryloxy; haloalkylcarbonyl; aryloxy; haloalkoxycarbonyl; alkylthiocarbonyl; arylthiocarbonyl; acyloxymethoxycarbonyl; cycloalkyl; bicycloalkyl; aryl; acyl; benzoyl; alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxy, haloalkyl, cyano, nitro, carboxyl derivatives, amino, alkoxy, thio, alkylthio, sulfonyl, aryl, aralkyl, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethyl, sulfonyl, alkylsulfonyl, haloalkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, fused monocyclic heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino,

- 807 -

amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles; monocyclic heterocycles; monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, aryloxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, aryl, fused aryl; monocyclic and bicyclic heterocyclicalkyls;  $-SO_2R^{10}$  wherein  $R^{10}$  is selected from the group consisting of alkyl, aryl and monocyclic heterocycles, all optionally substituted with one or more substituent selected from the group consisting of halogen, haloalkyl, alkyl, alkoxy, cyano, nitro, amino, acylamino, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, trifluoromethylthio, trifluoroalkoxy, trifluoromethylsulfonyl, aryl, aryloxy, thio, alkylthio, and monocyclic heterocycles; and




or  $NR^7$  and  $R^8$  taken together form a 4-12 membered mononitrogen containing monocyclic or bicyclic ring optionally substituted with one or more substituent selected from lower alkyl, carboxyl derivatives, aryl or hydroxy and wherein said ring optionally contains a heteroatom selected from the group consisting of O, N and S;

$R^5$  is selected from the group consisting of H, alkyl, alkenyl, alkynyl, benzyl, and phenethyl;



- 808 -

or




wherein Y<sup>2</sup> is selected from the group consisting of alkyl; cycloalkyl; bicycloalkyl; aryl; monocyclic heterocycles; alkyl optionally substituted with aryl which can also be optionally substituted with one or more substituent selected from halo, haloalkyl, alkyl, nitro, hydroxy, alkoxy, aryloxy, aryl, or fused aryl; aryl optionally substituted with one or more substituent selected from halo, haloalkyl, hydroxy, alkoxy, aryloxy, aryl, fused aryl, nitro, methylenedioxy, ethylenedioxy, or alkyl; alkynyl; alkenyl; -S-R<sup>9</sup> and -O-R<sup>9</sup> wherein R<sup>9</sup> is selected from the group consisting of H; alkyl; aralkyl; aryl; alkenyl; and alkynyl; or R<sup>9</sup> taken together with R<sup>7</sup> forms a 4-12 membered mononitrogen and monosulfur or monooxygen containing heterocyclic ring optionally substituted with lower alkyl, hydroxy, keto, phenyl, carboxyl or carboxyl ester, and fused phenyl; or R<sup>9</sup> taken together with R<sup>7</sup> is thiazole; oxazole; benzoxazole; or benzothiazole; and

R<sup>5</sup> and R<sup>7</sup> are as defined above;

or Y<sup>2</sup> (when Y<sup>2</sup> is carbon) taken together with R<sup>7</sup> forms a 4-12 membered mononitrogen containing ring optionally substituted with alkyl, aryl or hydroxy;

- 809 -


or A is



where  $\text{R}^2$  and  $\text{R}^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, phenyl, or carboxyl derivatives; and  $\text{R}^8$  is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, or acyloxymethoxycarbonyl; and

$\text{R}^5$  is defined as above

or A is



where  $\text{R}^2$  and  $\text{R}^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with hydroxy, keto, phenyl, or alkyl; and

$\text{R}^8$  are both selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl and acyloxymethoxycarbonyl;

- 810 -

Z<sup>1</sup> is one or more substituent s lected from th  
gr up c nsisting of H; alkyl; hydr xy; alk xy;  
aryloxy; halogen; haloalkyl; haloalkoxy; nitro;  
amino; alkylamino; acylamino; dialkylamino; cyano;  
alkylthio; alkylsulfonyl; carboxyl derivatives;  
trihaloacetamide; acetamide; aryl; fused aryl;  
cycloalkyl; thio; monocyclic heterocycles; fused  
monocyclic heterocycles; and A, wherein A is  
defined above;

V is selected from the group consisting of -N-(R<sup>6</sup>)-  
wherein R<sup>6</sup> is selected from the group consisting of  
H; lower alkyl; cycloalkyl; aralkyl; aryl; and  
monocyclic heterocycles; or R<sup>6</sup> taken together with  
Y, forms a 4-12 membered mononitrogen containing  
ring;

Y, Y<sup>3</sup>, Z and Z<sup>3</sup> are independently selected from the  
group consisting of hydrogen; alkyl; aryl; and  
cycloalkyl; or Y and Z taken together form a  
cycloalkyl; or Y<sup>3</sup> and Z<sup>3</sup> taken together form a  
cycloalkyl;

n is an integer 1, 2, or 3;

t is an integer 0, 1, or 2;

p is an integer 0, 1, 2, or 3;

R is X-R<sup>3</sup> wherein X is selected from the group  
consisting of O, S and NR<sup>4</sup>, wherein R<sup>3</sup> and R<sup>4</sup> are  
independently selected from the group consisting  
of hydrogen; alkyl; alkenyl; alkynyl; haloalkyl;  
aryl; arylalkyl; sugars; steroids;  
polyalkylethers; alkylamid ; alkyl N,N-  
dialkylamido; pivaloyloxymethyl; and in the case

- 811 -

of the free acid, all pharmaceutically acceptable salts thereof;

R<sup>1</sup> is selected from the group consisting of hydrogen; alkyl; alkenyl; alkynyl; aryl; carboxyl derivatives; haloalkyl; monocyclic heterocycles; monocyclic heterocycles optionally substituted with alkyl, halogen, haloalkyl, cyano, hydroxy, aryl, fused aryl, nitro, alkoxy, aryloxy, alkylsulfonyl, arylsulfonyl, sulfonamide, thio, alkylthio, carboxyl derivatives, amino, amido;

alkyl optionally substituted with one or more of halo, haloalkyl, hydroxy, alkoxy, aryloxy, thio, alkylthio, alkynyl, alkenyl, alkyl, arylthio, alkylsulfoxide, alkylsulfonyl, arylsulfoxide, arylsulfonyl, cyano, nitro, amino, alkylamino, dialkylamino, alkylsulfonamide, arylsulfonamide, acylamide, carboxyl derivatives, sulfonamide, sulfonic acid, phosphonic acid derivatives, phosphinic acid derivatives, aryl, arylthio, arylsulfoxide, or arylsulfone all optionally substituted on the aryl ring with halo, alkyl, haloalkyl, cyano, nitro, hydroxy, carboxyl derivatives, alkoxy, aryloxy, amino, alkylamino, dialkylamino, amido, aryl, fused aryl, monocyclic heterocycles; and fused monocyclic heterocycles, monocyclic heterocyclicthio, monocyclic heterocyclicsulfoxide, and monocyclic heterocyclic sulfone, which can be optionally substituted with halo, haloalkyl, nitro, hydroxy, alkoxy, fused aryl, or alkyl;

alkylcarbonyl, haloalkylcarbonyl, and arylcarbonyl;

- 812 -

aryl optionally substituted in one or more positions with halo, hal alkyl, alkyl, alkoxy, aryloxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, acyloxy, carboxyl derivatives, carboxyalkoxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycles; and



and provided that taken together with the nitrogen, R<sup>7</sup> and R<sup>8</sup> comprise an amino acid;


and

R<sup>11</sup> is selected from the group consisting of H, alkyl, aralkyl, alkenyl, alkynyl, haloalkyl or haloalkynyl or R<sup>11</sup> taken together with Y forms a 4-12 membered mononitrogen containing ring; and

a pharmaceutically acceptable carrier.

22. A pharmaceutical composition according to Claim 21 wherein

A is



wherein Y<sup>1</sup> is selected from the group consisting of N-R<sup>2</sup>, O, and S;

- 813 -

$R^2$  is selected from the group consisting of H, alkyl, aryl, substituted alkyl, hydroxy, alkoxy, alkylcarbonyl, cyano, nitro, amino and monocyclic heterocycles, arylcarbonyl, alkoxycarbonyl, aryloxy, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl; or  $R^2$  taken together with  $R^7$  forms a 4-12 membered ring;

$R^5$ ,  $R^7$ ,  $R^8$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, bicycloalkyl substituted alkyl, arylalkyl, aryloxy, hydroxy, alkoxy, amino, alkylamino, arylamino, amido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl, substituted phenyl, arylacyl, monocyclic and bicyclic heterocycles, monocyclic and bicyclic heterocyclicalkyl and  $-SO_2R^{10}$  wherein  $R^{10}$  is selected from the group consisting of alkyl, amino and aryl which are all optionally substituted with one or more substituent selected from the group consisting of acylamino, amino, carbonyl, cyano, nitro, alkoxy, halo, alkyl, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, aryloxy, thio, trifluoromethylthio, trifluoroalkoxy, and trifluoromethylsulfonyl or  $-NR^7$  and  $R^8$  taken together form a 4-12 membered ring wherein said ring optionally contains a heteroatom selected from the group consisting of O, N, and S wherein said ring is optionally substituted.

- 814 -

23. A pharmaceutical composition according to Claim 22 wherein

V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0 or 1; and

p is 0, 1 or 2.

24. A pharmaceutical composition according to Claim 23 wherein the compound is selected from the group consisting of

(±)ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±) $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

(±)ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

(±)ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoate;

- 815 -

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-naphthalen-1-yl]carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)-amino]-naphthalen-1-yl]carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

(±)ethyl β-[[1-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]-1-cyclopropyl]carbonyl]amino]pyridine-3-propanoate;

(±)β-[[1-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]cycloprop-1-yl]carbonyl]amino]pyridine-3-propanoic acid;

(±)ethyl β-[[2-[[[3-[[imino[(phenylmethyl)amino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[[imino[(phenylmethyl)amino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

3S-[[2-[[[3-(aminocarbonylamino)phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-(aminocarbonylamino)phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;



- 816 -

$\beta$ S-[[2-[[[3-[(aminoiminomethyl)amino]-2,5,6-trifluorophenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

ethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propanoic acid;

ethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propanoate;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-naphthalene-1-carboxylic acid;

- 817 -

methyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amin ]phenyl]carbonyl]amino]acetyl]amino]-  
naphthalene-1-carboxylate;

( $\pm$ ) 3-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]-2-oxopyrrolidine-1-  
propanoic acid;

3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoate;

3S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-pentynoic acid;

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

- 818 -

(±) ethyl β-[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±) ethyl β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]furan-2-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-furan-3-propanoic acid;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-amino]acetyl]amino]pentanedioic acid;

bismethyl ester 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]-amino]acetyl]amino]pentanedioate;

(±) hydrogen methyl 3-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]pentanedioate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-furan-2-propanoic acid;

ethyl (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]furan-2-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-naphthalene-2-propanoic acid;

(±) methyl β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-thiophene-3-propanoate;

- 819 -

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amin ]-  
4-pentynoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
thiophene-3-propanoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
carboxybutyl]thio]benzoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
carboxybutyl]sulfonyl]benzoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
thiophene-2-propanoic acid;

(±) methyl 2-[3-[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
carboxybutyl]thio]benzoate;

(±) methyl 3-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
5-[(4-methylphenyl)thio]pentanoate;

(±) methyl 3-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
4-[[4-methylphenyl)sulfonyl]amino]butanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-[[4-  
methylphenyl)sulfonyl]amino]butanoic acid;

- 820 -

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)thio]pentanoic acid;

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)sulfonyl]pentanoic acid;

3S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-(phenylthio)butanoic acid;

3R-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

2-[[2S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2-(carboxymethyl)ethyl]sulfonyl]benzoic acid;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

2-[[2S-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-2-(carboxymethyl)ethyl]thio]benzoic acid;

- 821 -

(±) thyl β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]methylamino]acetyl]-amino]-pyridine-3-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl) amino]phenyl]-carbonyl]methylamino]acetyl]amino]pyridine-3-propanoic acid;

(±) ethyl β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]-1-oxopropyl]-amino]pyridine-3-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl) amino]-phenyl]carbonyl]amino]-1-oxopropyl]-amino]pyridine-3-propanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl) amino]-4-methylphenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

(±) β-[[2-[[[3-[[[aminoiminomethyl) amino]-methyl]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

3S-[[2-[[[3-[(aminoiminomethyl) amino]-phenyl]carbonyl]amino]acetyl]amino]-4-hydroxybutanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl) amino]-phenyl]carbonyl]amino]acetyl]amino]-2-hydroxy-5-methylbenzenepropanoic acid;

- 822 -

(±) 3-[[2-[[[3-[(aminoiminomethyl)amin ]-phenyl]carbonyl]amino]acetyl]amino]-4-[(2-hydroxyethyl)amino]-4-oxobutanoic acid;

2S-[[2-[[[3-[[aminoiminomethyl]amino]phenyl]-carbonyl]amino]acetyl]amino]-3-carboxypropyl 2-aminobenzoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-1,4-benzodioxin-6-propanoic acid;

N-[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]-β-alanine, ethyl ester;

N-[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]-β-alanine;

(±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-quinoline-3-propanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-quinoline-3-propanoic acid;

N-[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]-β-alanine, ethyl ester;

3-[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]amino]propanoic acid;

(±) ethyl β-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

- 823 -

(±)  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(phenylamino)carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(phenylamino)carbonyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-  
propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminothioxomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;



- 824 -

ethyl  $\beta$ -[[2-[[[3-[(amin thioxomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-3-[[[(phenylamino)carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-3-[[[(phenylamino)carbonyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoic acid;

$\beta$ -[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]pyridine-  
3-propanoic acid;

- 825 -

$\beta$ -[[2-[[[3-[[[(3-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(2-carboxyethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(2-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(1-naphthalenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
chlorophenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
chlorophenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-chlorophenyl]-  
carbonyl]amino]acetyl]amino]benzenepropanoic acid;

ethyl  $\beta$ -[2-[[[3-[[amino(aminocarbonyl)-  
imino]methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(aminocarbonyl)imino]-  
methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzene-  
propanoic acid;

- 826 -

[(dimethylamino)carbonyl]methyl  $\beta$ -  
 [[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
 carbonyl]amino]acetyl]amino]-3,5-  
 dichlorobenzenepropanoate;

1,1-dimethylethyl 3,5-dichloro- $\beta$ -[[2-[[[3[[[(ethoxy-  
 carbonyl)amino][(ethoxycarbonyl)imino]methyl]amino]-  
 phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(ethoxycarbonyl)-  
 amino][(ethoxycarbonyl)imino]methyl]-  
 amino]-phenyl]carbonyl]amino]acetyl]-  
 amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
 chlorophenyl]carbonyl]amino]acetyl]amino]-  
 3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
 chlorophenyl]carbonyl]amino]acetyl]amino]-  
 3,5-dichlorobenzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]-  
 methyl]amino]phenyl]carbonyl]amino]acetyl]-  
 amino]-4-pentynoate;

3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]-  
 methyl]amino]phenyl]carbonyl]amino]-  
 acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)-  
 amino]-4-chlorophenyl]carbonyl]amino]-  
 acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-4-  
 chlorophenyl]carbonyl]amino]acetyl]-  
 amino]-4-pentynoic acid;

- 827 -

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoic acid;

- 828 -

- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3-  
bromobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3-  
bromobenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
bromobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-  
bromobenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dimethylbenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dimethylbenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dimethoxybenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dimethoxybenzenepropanoic acid;
- (±) (2,2-dimethyl-1-oxopropoxy)methyl  $\beta$ -[[2-[[[3-  
[(aminoiminomethyl)amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzenepropanoate;

- 829 -

- (±)  $\beta$ -[[2-[[[3-[[[(aminocarbonyl)imino)-methylamino)methyl]amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,4-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-fluoro-5-(trifluoromethyl)benzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-fluorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-methylbenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]-amino]-3,5-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-chlorobenzenepropanoic acid;

- 830 -

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-chlorobenzenepropanoic acid;

(±) [2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethyl]  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-iodobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2-hydroxy-4-methoxybenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-hydroxy-4-methoxybenzofuran-6-propanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichloro-2-hydroxybenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2-hydroxy-5-nitrobenzenepropanoic acid;

- 831 -

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-cyclohexanepropanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-cyclohexanepropanoate;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichloro-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-chloro-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-chloro-2-hydroxybenzenepropanoic acid;



- 832 -

(±) 5-amino-β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromopyridine-3-propanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromopyridine-3-propanoate;

3,5-dichloro-β-[[2-[[[3-[[[(ethoxycarbonyl)amino]-thioxomethyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

3,5-dichloro-β-[[2-[[[3-[[[(ethoxycarbonyl)amino]-iminomethyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino][1,1'-biphenyl]-3-propanoic acid;

1,1-dimethylethyl β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino][1,1'-biphenyl]-3-propanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino][pyrimidine-5-propanoic acid;

1,1-dimethylethyl β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-[pyrimidine-5-propanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-methylthiophene-2-propanoic acid;

- 833 -

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3-methylthiophene-2-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-(methylthio)benzenepropanoic acid;

1,1-dimethylethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3-(methylthio)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-6-methylpyridine-2-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-(methylsulfonyl)benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-diethoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-diethoxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromothiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromothiophene-2-propanoate;

- 834 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl-carbonyl]amino]acetyl]amino]-5-chlorothiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl-carbonyl]amino]acetyl]amino]-5-chlorothiophene-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-1H-pyrazole-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-1H-pyrazole-3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-methylthiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-methylthiophene-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2,3,5-trichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2,3,5-trichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-benzenepropanoic acid;

- 835 -

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-6-chloro-  
1,3-benzodioxole-5-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-6-chloro-  
1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzofuran-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzofuran-2-propanoate;

- 836 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-(carboxymethoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-(carboxymethoxy)benzenepropanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4,4,4-trifluorobutanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4,4,4-trifluorobutanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-4,5-dimethoxybenzenepropanoic acid;

ethyl( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-4,5-dimethoxybenzenepropanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-methylpentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-methylpentanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-phenyl]carbonyl]amino]acetyl]amino]-pentanoic acid;

- 837 -

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pentanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
5-bromo-3-chloro-2-hydroxybenzene-  
propanoic acid;

$\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

- 838 -

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]pyridine-3-  
propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

- 839 -

3-[[2-[[[3-[[[(3,5-dichlor phenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]butanoic acid;

ethyl 3-[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]butanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-bis(1-methyl-  
ethoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-bis(1-methyl-  
ethoxy)benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dibromo-4-  
hydroxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dibromo-4-  
hydroxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dichloro-4-  
hydroxybenzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dichloro-4-  
hydroxybenzenepropanoate;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-  
hydroxyphenyl]carbonyl]amino]acetyl]amino]-  
3,5-dichlorobenzenepropanoic acid;



- 840 -

ethyl  $\beta$ -[[2-[[[5-[(amin iminomethyl)amino]-2-hydr xyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(phenoxyamino)carbonyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(phenoxyamino)carbonyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylamino)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylamino)amino]carbonyl]-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-[(3,5-dichlorophenyl)amino]-5-oxopentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-[(3,5-dichlorophenyl)amino]-5-oxopentanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

- 841 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-(acetylamino)-5-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-(acetylamino)-5-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(methylamino)(methylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

- 842 -

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-  
[[ (thylamino) (m thylimino) methyl] amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
benzenepropanoic acid;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(1-  
methylethyl) amino] (methylimino) methyl] amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
benzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[[ (ethylamino)-  
(methylimino) methyl] amino] phenyl] carbonyl]-  
amino] acetyl] amino]-4-fluorobenzene-  
propanoic acid;

(±) 4-fluoro- $\beta$ -[[2-[[[3-[[[(4-  
pyridinylmethyl) amino] (methylimino) methyl]-  
amino] phenyl] carbonyl] amino] acetyl] amino]-  
benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl) amino] phenyl]-  
carbonyl] amino] acetyl] amino]-4-  
fluorobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl) amino]-  
phenyl] carbonyl] amino] acetyl] amino]-  
1H-imidazole-2-propanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl) amino] phenyl]-  
carbonyl] amino] acetyl] amino]-2,3,4,6-  
tetrafluorobenzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino] phenyl] carbonyl] amino] acetyl] amino]-  
5-bromothiophene-2-propanoic acid;

- 843 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-2-  
mercaptobenzenepropanoic acid;

and

( $\pm$ )  $\beta$ -[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-5-  
chloro-2-mercaptobenzenepropanoic acid.

25. A pharmaceutical composition according to Claim 23  
wherein  $Y^1$  is  $N-R^2$  and  $R^2$  is cyano.

26. A pharmaceutical composition according to Claim 25  
wherein the compound is selected from the group  
consisting of

phenylmethyl  $\beta$ -[[2-[[[3-[[[(cyanoimino)phenylmethyl-  
amino)methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[[[(cyanoimino)-  
methylanino)methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[[[(cyanoimino)-  
(amino)methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(cyanoimino)(ethylamino)methyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(cyanoimino)[(phenylmethyl)-  
amino)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]benz nepropanoic acid;

- 844 -

$\beta$ -[[2-[[[3-[(cyan imino)(methylamino)-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[amino(cyanoimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

$\beta$ -[[2-[[[3-[(cyanoimino)(ethylamino)-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

ethyl 3S-[[2-[[[3-[(cyanoimino)-(methylamino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(cyanoimino)(methylamino)-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

ethyl  $\beta$ -[[2-[[[3-[(cyanoimino)[2-pyridinylmethyl]-amino]methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)[2-pyridinylmethyl]-amino]methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(cyanoimino)[3-pyridinylmethyl]amino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)[3-pyridinylmethyl]-amino]methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoic acid;

- 845 -

ethyl 3S-[[2-[[[3-[[[(4-(aminosulfonyl)-  
phenylmethyl]amino)(cyan imino)methyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoate;

3S-[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl)-  
amino](cyanoimino)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-4-pentynoic acid;

ethyl  $\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoic acid;

( $\pm$ ) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)-  
(methylamino)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoate;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)(methylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;


and

ethyl 3-[[2-[[[3-[[amino(cyanoimino)methyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate.

- 846 -

27. A pharmaceutical composition according to Claim 21 wherein

A is



wherein Y<sup>1</sup> is N-R<sup>2</sup>; R<sup>2</sup> taken together with R<sup>7</sup> forms a 4-12 membered ring; and R<sup>8</sup> is selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, bicycloalkyl substituted alkyl, arylalkyl, aryloxy, hydroxy, alkoxy, amino, alkylamino, arylamino, amido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl, substituted phenyl, arylacyl, monocyclic and bicyclic heterocycles, monocyclic and bicyclic heterocyclicalkyl and -SO<sub>2</sub>R<sup>10</sup> wherein R<sup>10</sup> is selected from the group consisting of alkyl, amino and aryl which are all optionally substituted with one or more substituent selected from the group consisting of acylamino, amino, carbonyl cyano, nitro, alkoxy, halo, alkyl, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, aryloxy, thio, trifluoromethylthio, trifluoroalkoxy, and trifluoromethylsulfonyl or -NR<sup>7</sup> and R<sup>8</sup> taken together form a 4-12 membered ring wherein said ring optionally contains a heteroatom selected from the group consisting of O, N, and S wherein said ring is optionally substituted.

- 847 -

28. A pharmaceutical composition according to Claim 27 wherein

V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0; and

p is 1.

29. A pharmaceutical composition according to Claim 28 wherein the compound is selected from the group consisting of

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;



- 848 -

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

(±) 3,5-dibromo- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

(±) 3-bromo-5-chloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

(±) [2-[2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) [2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) 3-bromo-5-chloro-2-hydroxy- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

- 849 -

ethyl (±) 3-bromo-5-chloro-2-hydroxy-β-  
[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

(±) 3,5-dichloro-2-hydroxy-β-[[2-  
[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

(±) 3,5-dichloro-β-[[2-[[[3-[(4,5-dihydro-  
1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]-  
amino]acetyl]amino]-2-hydroxybenzenepropanoic acid;

(±) 3-bromo-5-chloro-β-[[2-[[[3-[(4,5-dihydro-  
1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]-  
acetyl]amino]-2-hydroxybenzenepropanoic acid;

β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

(±) ethyl 3,5-dichloro-β-[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

(±) 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-  
dimethylpyrimidin-2-yl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoic acid;

- 850 -


ethyl ( $\pm$ ) 3,5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

( $\pm$ ) 3-bromo-5-dichloro-2-hydroxy-  
 $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-  
dimethylpyrimidin-2-yl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]benzene-  
propanoic acid;

and

ethyl( $\pm$ ) 3-bromo-5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-  
[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-  
yl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate.

30. A pharmaceutical composition according to Claim 21 wherein



wherein  $\text{Y}^2$  is selected from the group consisting of lower alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, monocyclic heterocycles,  $-\text{S}-\text{R}^9$  and  $-\text{O}-\text{R}^9$  wherein  $\text{R}^9$  is selected from the group consisting of H, alkyl, substituted alkyl, phenyl, substituted phenyl and monocyclic heterocycles or  $\text{R}^9$  taken together with  $\text{R}^7$  forms a 4-12 membered ring; or

- 851 -

Y<sup>2</sup> taken together with R<sup>7</sup> forms a 4-12 membered ring which is optionally substituted.

31. A pharmaceutical composition according to Claim 30 wherein

Y<sup>2</sup> taken together with R<sup>7</sup> forms a 4-12 membered ring which is optionally substituted.

32. A pharmaceutical composition according to Claim 31 wherein

V is -N(R<sup>6</sup>)- wherein R<sup>6</sup> is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0 or 1; and

p is 1.

33. A pharmaceutical composition according to Claim 32 wherein the compound is selected from the group consisting of

(±) ethyl β-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±) β-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

(±) ethyl β-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoate;

- 852 -

- (±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;
- $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoate;
- (±)  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoic acid;
- (±)  $\beta$ -[[2-[[[3,5-bis[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;
- $\beta$ S-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

- 853 -

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

(±) 3,5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

and

ethyl(±) 3,5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate.

- 854 -

34. A pharmaceutical composition according to Claim 30 wherein

$Y^2$  is selected from the group consisting of lower alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl and monocyclic heterocycles.

35. A pharmaceutical composition according to Claim 34 wherein

V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H and lower alkyl;

n is 1;

t is 0; and

p is 1.

36. A pharmaceutical composition according to Claim 35 wherein the compound is selected from the group consisting of

(±) ethyl  $\beta$ -[[2-[[[3-[(iminophenylmethyl)-amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ S-[[2-[[[3-[[imino(1-pyrrolidinyl)methyl]-amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

(±)  $\beta$ -[[2-[[[3-[(iminophenylmethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

- 855 -

3,5-dichloro- $\beta$ -[[2-[[[3-[[imino(1-piperidinyl)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

and

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[imino(1-piperidinyl)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate.

37. A pharmaceutical composition according to Claim 30 wherein  $Y^2$  is  $-S-R^9$  or  $-O-R^9$  wherein  $R^9$  is selected from the group consisting of H, alkyl, substituted alkyl, phenyl, substituted phenyl and monocyclic heterocycles or  $R^9$  taken together with  $R^7$  forms a 4-12 membered ring.
38. A pharmaceutical composition according to Claim 37 wherein
- V is  $-N(R^6)-$  wherein  $R^6$  is selected from the group consisting of H, lower alkyl, substituted alkyl, cycloalkyl, aryl, substituted aryl, monocyclic heterocycles and benzyl;
- n is 0;
- t is 0; and
- p is 1 or 2.
39. A pharmaceutical composition according to Claim 38 wherein the compound is selected from the group consisting of



- 856 -

ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydrothiazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(4,5-dihydrothiazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[benzoxazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;


ethyl  $\beta$ -[[2-[[[3-[[benzoxazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate.

$\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;



and

ethyl  $\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate.





40. A pharmaceutical composition according to Claim 21 wherein the compound is selected from the group consisting of








- 857 -








- 858 -







- 859 -







- 860 -







- 861 -



- 862 -







- 863 -














- 864 -







- 865 -







- 866 -






- 867 -






- 868 -







- 869 -



- 870 -






- 871 -










- 872 -






- 873 -






- 874 -






- 875 -






- 876 -






- 877 -



- 878 -







- 879 -








- 880 -



- 881 -




41. A method for treating conditions mediated by the  $\alpha_v\beta_3$  integrin in a mammal in need of such treatment comprising administering an effective  $\alpha_v\beta_3$  inhibiting amount of a compound of the formula



or a pharmaceutically acceptable salt thereof,  
wherein

A is



- 882 -

wherein  $Y^1$  is selected from the group consisting of  $N-R^2$ , O, and S;

$R^2$  is selected from the group consisting of H; alkyl; aryl; hydroxy; alkoxy; cyano; nitro; amino; alkenyl; alkynyl; amido; alkylcarbonyl; arylcarbonyl; alkoxycarbonyl; aryloxycarbonyl; haloalkylcarbonyl; haloalkoxycarbonyl; alkylthiocarbonyl; arylthiocarbonyl; acyloxymethoxycarbonyl; alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxyl, haloalkyl, cyano, nitro, carboxyl, amino, alkoxy, aryl or aryl optionally substituted with one or more halogen, haloalkyl, lower alkyl, alkoxy, cyano, alkylsulfonyl, alkylthio, nitro, carboxyl, amino, hydroxyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, hydroxy, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, cyano, nitro, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, carboxyl derivatives, amino, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycle; monocyclic heterocycles; and monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, aryl or fused aryl; or

$R^2$  taken together with  $R^1$  forms a 4-12 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy,

- 883 -

k to, alkyl, halo, phenyl, amino, carboxyl or carboxyl ester, and fused phenyl;


or R<sup>2</sup> taken together with R<sup>7</sup> forms a 5 membered heteroaromatic ring optionally substituted with one or more substituent selected from lower alkyl, phenyl and hydroxy;

or R<sup>2</sup> taken together with R<sup>7</sup> forms a 5 membered heteroaromatic ring fused with a phenyl group;

R<sup>7</sup> (when not taken together with R<sup>2</sup>) and R<sup>8</sup> are independently selected from the group consisting of H; alkyl; alkenyl; alkynyl; aralkyl; amino; alkylamino; hydroxy; alkoxy; arylamino; amido, alkylcarbonyl, arylcarbonyl; alkoxycarbonyl; aryloxy; haloalkylcarbonyl; aryloxy; haloalkoxycarbonyl; alkylthiocarbonyl; arylthiocarbonyl; acyloxymethoxycarbonyl; cycloalkyl; bicycloalkyl; aryl; acyl; benzoyl; alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxy, haloalkyl, cyano, nitro, carboxyl derivatives, amino, alkoxy, thio, alkylthio, sulfonyl, aryl, aralkyl, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethyl, sulfonyl, alkylsulfonyl, haloalkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, fused monocyclic heterocycles; aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy,

- 884 -

methylenedioxy, thylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles; monocyclic heterocycles; monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, aryloxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, aryl, fused aryl; monocyclic and bicyclic heterocyclicalkyls;  $-SO_2R^{10}$  wherein  $R^{10}$  is selected from the group consisting of alkyl, aryl and monocyclic heterocycles, all optionally substituted with one or more substituent selected from the group consisting of halogen, haloalkyl, alkyl, alkoxy, cyano, nitro, amino, acylamino, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, trifluoromethylthio, trifluoroalkoxy, trifluoromethylsulfonyl, aryl, aryloxy, thio, alkylthio, and monocyclic heterocycles; and




or  $NR^7$  and  $R^8$  taken together form a 4-12 membered mononitrogen containing monocyclic or bicyclic ring optionally substituted with one or more substituent selected from lower alkyl, carboxyl derivatives, aryl or hydroxy and wherein said ring optionally contains a heteroatom selected from the group consisting of O, N and S;

- 885 -

R<sup>5</sup> is selected from the group consisting of H, alkyl, alkenyl, alkynyl, benzyl, and phenethyl;

or




wherein Y<sup>2</sup> is selected from the group consisting of alkyl; cycloalkyl; bicycloalkyl; aryl; monocyclic heterocycles; alkyl optionally substituted with aryl which can also be optionally substituted with one or more substituent selected from halo, haloalkyl, alkyl, nitro, hydroxy, alkoxy, aryloxy, aryl, or fused aryl; aryl optionally substituted with one or more substituent selected from halo, haloalkyl, hydroxy, alkoxy, aryloxy, aryl, fused aryl, nitro, methylenedioxy, ethylenedioxy, or alkyl; alkynyl; alkenyl; -S-R<sup>9</sup> and -O-R<sup>9</sup> wherein R<sup>9</sup> is selected from the group consisting of H; alkyl; aralkyl; aryl; alkenyl; and alkynyl; or R<sup>9</sup> taken together with R<sup>7</sup> forms a 4-12 membered mononitrogen and monosulfur or monooxygen containing heterocyclic ring optionally substituted with lower alkyl, hydroxy, keto, phenyl, carboxyl or carboxyl ester, and fused phenyl; or R<sup>9</sup> taken together with R<sup>7</sup> is thiazole; oxazole; benzoxazole; or benzothiazole; and

R<sup>5</sup> and R<sup>7</sup> are as defined above;

or Y<sup>2</sup> (when Y<sup>2</sup> is carbon) taken together with R<sup>7</sup> forms a 4-12 membered mononitrogen containing ring optionally substituted with alkyl, aryl or hydroxy;

- 886 -


or A is



where  $\text{R}^2$  and  $\text{R}^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, phenyl, or carboxyl derivatives; and  $\text{R}^8$  is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, or acyloxymethoxycarbonyl; and

$\text{R}^5$  is defined as above

or A is



where  $\text{R}^2$  and  $\text{R}^7$  taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with hydroxy, keto, phenyl, or alkyl; and

$\text{R}^8$  are both selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl and acyl xymethoxycarbonyl;

- 887 -

Z<sup>1</sup> is one or more substituent selected from the group consisting of H; alkyl; hydroxy; alkoxy; aryloxy; halogen; haloalkyl; haloalkoxy; nitro; amino; alkylamino; acylamino; dialkylamino; cyano; alkylthio; alkylsulfonyl; carboxyl derivatives; trihaloacetamide; acetamide; aryl; fused aryl; cycloalkyl; thio; monocyclic heterocycles; fused monocyclic heterocycles; and A, wherein A is defined above;

V is selected from the group consisting of -N-(R<sup>6</sup>)- wherein R<sup>6</sup> is selected from the group consisting of H; lower alkyl; cycloalkyl; aralkyl; aryl; and monocyclic heterocycles; or R<sup>6</sup> taken together with Y, forms a 4-12 membered mononitrogen containing ring;

Y, Y<sup>3</sup>, Z and Z<sup>3</sup> are independently selected from the group consisting of hydrogen; alkyl; aryl; and cycloalkyl; or Y and Z taken together form a cycloalkyl; or Y<sup>3</sup> and Z<sup>3</sup> taken together form a cycloalkyl;

n is an integer 1, 2, or 3;

t is an integer 0, 1, or 2;

p is an integer 0, 1, 2, or 3;

R is X-R<sup>3</sup> wherein X is selected from the group consisting of O, S and NR<sup>4</sup>, wherein R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydrogen; alkyl; alkenyl; alkynyl; haloalkyl; aryl; arylalkyl; sugars; steroids; polyalkylethers; alkylamido; alkyl N,N-dialkylamido; pivaloyloxymethyl; and in the case



- 888 -

of the free acid, all pharmaceutically acceptable salts thereof;


R<sup>1</sup> is selected from the group consisting of hydrogen; alkyl; alkenyl; alkynyl; aryl; carboxyl derivatives; haloalkyl; monocyclic heterocycles; monocyclic heterocycles optionally substituted with alkyl, halogen, haloalkyl, cyano, hydroxy, aryl, fused aryl, nitro, alkoxy, aryloxy, alkylsulfonyl, arylsulfonyl, sulfonamide, thio, alkylthio, carboxyl derivatives, amino, amido;

alkyl optionally substituted with one or more of halo, haloalkyl, hydroxy, alkoxy, aryloxy, thio, alkylthio, alkynyl, alkenyl, alkyl, arylthio, alkylsulfoxide, alkylsulfonyl, arylsulfoxide, arylsulfonyl, cyano, nitro, amino, alkylamino, dialkylamino, alkylsulfonamide, arylsulfonamide, acylamide, carboxyl derivatives, sulfonamide, sulfonic acid, phosphonic acid derivatives, phosphinic acid derivatives, aryl, arylthio, arylsulfoxide, or arylsulfone all optionally substituted on the aryl ring with halo, alkyl, haloalkyl, cyano, nitro, hydroxy, carboxyl derivatives, alkoxy, aryloxy, amino, alkylamino, dialkylamino, amido, aryl, fused aryl, monocyclic heterocycles; and fused monocyclic heterocycles, monocyclic heterocyclicthio, monocyclic heterocyclicsulfoxide, and monocyclic heterocyclic sulfone, which can be optionally substituted with halo, haloalkyl, nitro, hydroxy, alkoxy, fused aryl, or alkyl;

alkylcarbonyl, haloalkylcarbonyl, and arylcarbonyl;

- 889 -

aryl optionally substituted in one or more positions with halo, haloalkyl, alkyl, alkoxy, aryloxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, acyloxy, carboxyl derivatives, carboxyalkoxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycles; and



and provided that taken together with the nitrogen,  $\text{R}^7$  and  $\text{R}^8$  comprise an amino acid;

and

$\text{R}^{11}$  is selected from the group consisting of H, alkyl, aralkyl, alkenyl, alkynyl, haloalkyl or haloalkynyl or  $\text{R}^{11}$  taken together with Y forms a 4-12 membered mononitrogen containing ring.

42. A method according to claim 41 wherein the compound is selected from

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

- 890 -

- (±)β-[[2-[[[3-  
[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoic acid;
- (±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoate;
- (±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoic acid;
- (±)ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
naphthalen-1-yl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;
- (±)β-[[2-[[[3-[(aminoiminomethyl)-amino]-  
naphthalen-1-yl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;
- (±)ethyl β-[[1-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]-1-cyclopropyl]-  
carbonyl]amino]pyridine-3-propanoate;
- (±)β-[[1-[[[3[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]cycloprop-1-yl]carbonyl]-  
amino]pyridine-3-propanoic acid;
- (±)ethyl β-[[2-[[[3-[[imino[(phenylmethyl)amino]-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

- 891 -

(±)  $\beta$ -[[2-[[[3-[[imino[(ph nylmethyl)amino]methyl]-amino]ph nyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

3S-[[2-[[[3-(aminocarbonylamino)phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-(aminocarbonylamino)phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoate;

$\beta$ S-[[2-[[[3-[(aminoiminomethyl)amino]-2,5,6-trifluorophenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

ethyl (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propanoic acid;

ethyl (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-[1,1'-biphenyl]-4-propan ate;

- 892 -

(±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)-phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-naphthalene-1-carboxylic acid;

methyl (±) β-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-naphthalene-1-carboxylate;

(±) 3-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]-2-oxopyrrolidine-1-propanoic acid;

3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[[[(phenylmethyl)amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(aminoiminomethyl)amin]phenyl]-carbonyl]amino]acetyl]amino]-4-pentynoic acid;

- 893 -

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]furan-2-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
furan-3-propanoic acid;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]pentanedioic acid;

bismethyl ester 3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]-amino]acetyl]amino]pentanedioate;

( $\pm$ ) hydrogen methyl 3-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pentanedioate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
furan-2-propanoic acid;

- 894 -

ethyl (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]furan-2-propanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-naphthalene-2-propanoic acid;

(±) methyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-thiophene-3-propanoate;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-thiophene-3-propanoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-carboxybutyl]thio]benzoic acid;

(±) 2-[3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-carboxybutyl]sulfonyl]benzoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-thiophene-2-propanoic acid;

(±) methyl 2-[[3-[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-carboxybutyl]thio]benzoate;

- 895 -

(±) methyl 3-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)thio]pentanoate;

(±) methyl 3-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-4-[[4-methylphenyl)sulfonyl]amino]butanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-[[4-methylphenyl)sulfonyl]amino]butanoic acid;

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)thio]pentanoic acid;

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-[(4-methylphenyl)sulfonyl]pentanoic acid;

3S-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-(phenylthio)butanoic acid;

3R-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3R-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

2-[[2S-[[2-[[[3-[(aminoiminomethyl)-amino]phenyl]carbonyl]amino]acetyl]amino]-2-(carboxymethyl)ethyl)sulfonyl]benzoic acid;



- 896 -

3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-pentynoate;

2-[[2S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-2-  
(carboxymethyl)ethyl]thio]benzoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]methylamino]acetyl]-  
amino]-pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]methylamino]acetyl]amino]pyridine-  
3-propanoic acid;

(±)ethyl β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]-1-oxopropyl]-  
amino]pyridine-3-propanoate;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]-1-oxopropyl]-  
amino]pyridine-3-propanoic acid;

(±)β-[[2-[[[3-[(aminoiminomethyl)amino]-4-  
methylphenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

(±)β-[[2-[[[3-[[[aminoiminomethyl]amino]-  
methyl]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

- 897 -

3S-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
4-hydroxybutanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-2-  
hydroxy-5-methylbenzenepropanoic acid;

(±) 3-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-[(2-  
hydroxyethyl)amino]-4-oxobutanoic acid;

2S-[[2-[[[3-[[aminoiminomethyl]amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-  
carboxypropyl 2-aminobenzoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1,4-  
benzodioxin-6-propanoic acid;

N-[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]-β-alanine, ethyl ester;

N-[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]-β-alanine;

(±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
quinoline-3-propanoate;

β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
quinoline-3-propanoic acid;

- 898 -

N-[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
 $\beta$ -alanine, ethyl ester;

3-[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]  
amino]propanoic acid;

( $\pm$ )ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]  
acetyl]amino]pyridine-3-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylamino)carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylamino)carbonyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-(aminocarbonylamino)-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-(aminocarbonylamino)phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-3-  
propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

- 899 -

$\beta$ -[[2-[[[3-[[[(4-methylphenyl)sulfonyl]amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminothioxomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoic acid;

- 900 -

ethyl  $\beta$ -[[2-[[[3-3-[(phenylamino)carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-3-[(phenylamino)carbonyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-1,3-  
benzodioxole-5-propanoic acid;

$\beta$ -[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]pyridine-  
3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(3-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(2-carboxyethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(2-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

$\beta$ -[[2-[[[3-[[[(1-naphthalenylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-  
chlorophenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

- 901 -

$\beta$ -[[2-[[[3-[(amin iminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-chlorophenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

ethyl  $\beta$ -[2-[[[3-[[amino(aminocarbonyl)imino]methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(aminocarbonyl)imino]methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]-3,5-dichlorobenzene-propanoic acid;

[(dimethylamino)carbonyl]methyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

1,1-dimethylethyl 3,5-dichloro- $\beta$ -[[2-[[[3[[[(ethoxycarbonyl)amino][(ethoxycarbonyl)imino]methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(ethoxycarbonyl)amino][(ethoxycarbonyl)imino]methyl]-amino]-phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

- 902 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[[amino[(aminocarbonyl)imino]methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

ethyl 3S-[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[(aminoiminomethyl)amino]-4-chlorophenyl]carbonyl]amino]acetyl]amino]-4-pentynoic acid;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoic acid;

( $\pm$ ) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)carbonyl]amino]acetyl]amino]-3,4-dichlorobenzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

- 903 -

- (±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoate;
- (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2,5-dimethylbenzenepropanoic acid;
- (±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoate;
- (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-chlorobenzenepropanoic acid;
- (±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromobenzenepropanoate;
- (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-bromobenzenepropanoic acid;
- (±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromobenzenepropanoate;
- (±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-4-bromobenzenepropanoic acid;
- (±) ethyl β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethylbenzenepropanoate;



- 904 -

- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethylbenzenepropanoic acid;
- (±) ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethoxybenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dimethoxybenzenepropanoic acid;
- (±) (2,2-dimethyl-1-oxopropoxy)methyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[[[(aminocarbonyl)imino)-methylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminothioxomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,4-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3-fluoro-5-(trifluoromethyl)benzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-fluorobenzenepropanoic acid;

- 905 -

- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-methylbenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]-amino]-3,5-dibromobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-5-chlorobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoromethyl)phenyl]carbonyl]amino]acetyl]-amino]-3-bromo-5-chlorobenzenepropanoic acid;
- (±) [2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]-ethyl]  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-bromo-5-iodobenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2-hydroxy-4-methoxybenzenepropanoic acid;
- (±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-5-hydroxy-4-methoxybenzofuran-6-propanoic acid;

- 906 -

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-9H-fluorene-2-propanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-2-hydroxybenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-2-hydroxy-5-nitrobenzenepropanoic acid;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dibromo-2-hydroxybenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoic acid;

ethyl(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-5-bromo-2-hydroxybenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-cyclohexanepropanoic acid;

- 907 -

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
cyclohexanepropanoate;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
3,5-dichloro-2-hydroxybenzenepropanoate;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-5-  
chloro-2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-bromo-5-  
chloro-2-hydroxybenzenepropanoic acid;

(±) 5-amino-β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
2-hydroxybenzenepropanoic acid;

(±) β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
5-bromopyridine-3-propanoic acid;

ethyl(±) β-[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
5-bromopyridine-3-propanoate;

3,5-dichloro-β-[[2-[[[3-[[[(ethoxycarbonyl)amino]-  
thioxomethyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

3,5-dichloro-β-[[2-[[[3-[[[(ethoxycarbonyl)amino]-  
iminomethyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

- 908 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino][1,1'-  
biphenyl]-3-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino][1,1'-  
biphenyl]-3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino][pyrimidine-5-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
[pyrimidine-5-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]-  
amino]acetyl]amino]-3-methylthiophene-2-propanoic acid;

1,1-dimethylethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-3-  
methylthiophene-2-propanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-(methylthio)-  
benzenepropanoic acid;

1,1-dimethylethyl ( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
3-(methylthio)benzenepropanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-6-  
methylpyridine-2-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-(methylsulfonyl)-  
benzenepropanoic acid;

- 909 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-  
diethoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]-carbonyl]amino]acetyl]amino]-3,5-  
diethoxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
bromothiophene-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
bromothiophene-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-chlorothiophene-  
2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-chlorothiophene-  
2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1H-pyrazole-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-1H-pyrazole-  
3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-methylthiophene-  
2-propanoic acid;

- 910 -

thyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-m thylthi phen -  
2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2,3,5-trichloro-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2,3,5-trichloro-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-2(carboxymethoxy)-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-methoxy-1,3-  
benzodioxole-6-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-bromo-  
2-methoxybenzenepropanoate;

- 911 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-6-chloro-1,3-benzodioxole-5-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-6-chloro-1,3-benzodioxole-5-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-benzofuran-2-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-benzofuran-2-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-(carboxymethoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-phenyl]carbonyl]amino]acetyl]amino]-3-(carboxymethoxy)benzenepropanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4,4,4-trifluorobutanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4,4,4-trifluorobutanoate;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3-bromo-4,5-dimethoxybenzenepropanoic acid;



- 912 -

ethyl(±) β-[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3-bromo-4,5-  
dimethoxybenzenepropanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
methylpentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-4-  
methylpentanoate;

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pentanoate;

β-[2-[[[3-[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
5-bromo-3-chloro-2-hydroxybenzene-  
propanoic acid;

β-[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoic acid;

ethyl β-[2-[[[3-[[[(4-pyridinylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]pyridine-3-propanoate;

3,5-dichloro-β-[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

- 913 -

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(2-pyridinylmethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)amino]-  
carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;

ethyl 3-chloro- $\beta$ -[[2-[[[3-[[[(phenylmethyl)-  
amino]carbonyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1-phenylethyl)amino]carbonyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

- 914 -

$\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]amino]-carbonyl]amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(1H-benzimidazol-2-yl)-methyl]-amino]carbonyl]amino]phenyl]carbonyl]amino]-acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-amino]carbonyl]amino]phenyl]carbonyl]-amino]acetyl]amino]pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-amino]carbonyl]amino]phenyl]carbonyl]amino]-acetyl]amino]pyridine-3-propanoate;

3-[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-amino]carbonyl]amino]phenyl]carbonyl]-amino]acetyl]amino]butanoic acid;

ethyl 3-[[2-[[[3-[[[(3,5-dichlorophenyl)methyl]-amino]carbonyl]amino]phenyl]carbonyl]-amino]acetyl]amino]butanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-bis(1-methylethoxy)benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-bis(1-methylethoxy)benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dibromo-4-hydroxybenzenepropanoic acid;

- 915 -

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dibromo-4-hydroxybenzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-4-hydroxybenzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichloro-4-hydroxybenzenepropanoate;

$\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-hydroxyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[5-[(aminoiminomethyl)amino]-2-hydroxyphenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[(phenoxyamino)carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(phenoxyamino)carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[[[(phenylamino)amino]carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[[[(phenylamino)amino]carbonyl]amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

- 916 -

3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-[(3,5-  
dichlorophenyl)amino]-5-oxopentanoic acid;

ethyl 3-[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-5-[(3,5-  
dichlorophenyl)amino]-5-oxopentanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-  
phenyl]carbonyl]amino]acetyl]amino]pyridine-  
3-propanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-carboxy-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-  
dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]phenyl]-  
carbonyl]amino]acetyl]amino]-3,5-dichloro-  
benzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3,5-bis[(aminoiminomethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-3,5-dichloro-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-  
(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]-3,5-dichlorobenzenepropanoic acid;

- 917 -

ethyl  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]-5-(trifluoroacetyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-(acetyl)amino]-5-(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

ethyl  $\beta$ -[[2-[[[3-(acetyl)amino]-5-(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[(methylamino)(methyylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[(ethylamino)(methyylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[[[(1-methylethyl)amino](methyylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(ethylamino)(methyylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-4-fluorobenzenepropanoic acid;

( $\pm$ ) 4-fluoro- $\beta$ -[[2-[[[3-[[[(4-pyridinylmethyl)amino](methyylimino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

- 918 -

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]-carbonyl]amino]acetyl]amino]-4-fluorobenzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-1H-imidazole-2-propanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2,3,4,6-tetrafluorobenzenepropanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-bromothiophene-2-propanoic acid;

$\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-2-mercaptobenzenepropanoic acid;

( $\pm$ )  $\beta$ -[[2-[[[3-[(aminoiminomethyl)amino]phenyl]carbonyl]amino]acetyl]amino]-5-chloro-2-mercaptobenzenepropanoic acid;

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)phenylmethylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)methylamino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

phenylmethyl  $\beta$ -[[2-[[[3-[(cyanoimino)(amino)methyl]amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

- 919 -

$\beta$ -[[2-[[[3-[[ (cyanoimino) (ethylamino) methyl]-  
amino]ph nyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

$\beta$ -[[2-[[[3-[[ (cyanoimino) [(phenylmethyl)-  
amino]methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[[ (cyanoimino) (methylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

$\beta$ -[[2-[[[3-[[ amino (cyanoimino) methyl]amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoic acid;

$\beta$ -[[2-[[[3-[[ (cyanoimino) (ethylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]benzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[ (cyanoimino)-  
(methylamino) methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]-4-pentynoate;

3S-[[2-[[[3-[[ (cyanoimino) (methylamino)-  
methyl]amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoic acid;

ethyl  $\beta$ -[[2-[[[3-[[ (cyanoimino) [2-pyridinylmethyl)-  
amino]methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[[ (cyanoimino) [2-pyridinylmethyl)-  
amino]methyl]amino]phenyl]carbonyl]amino]-  
acetyl]amino]benzenepropanoic acid;



- 920 -

ethyl  $\beta$ -[[2-[[[3-[(cyanoimino)[3-pyridinylmethyl]amino]methyl]amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

$\beta$ -[[2-[[[3-[(cyanoimino)[3-pyridinylmethyl]-amino]methyl]amino]phenyl]carbonyl]amino]-acetyl]amino]benzenepropanoic acid;

ethyl 3S-[[2-[[[3-[[[(4-(aminosulfonyl)-phenylmethyl]amino)(cyanoimino)methyl]-amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoate;

3S-[[2-[[[3-[[[(4-(aminosulfonyl)phenylmethyl)-amino](cyanoimino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]-4-pentynoic acid;

ethyl  $\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

$\beta$ -[[2-[[[3-[[amino(cyanoimino)methyl]amino]phenyl]-carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

( $\pm$ ) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)-(methylamino)methyl]amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate;

( $\pm$ ) 3,5-dichloro- $\beta$ -[[2-[[[3-[(cyanoimino)(methylamino)-methyl]amino]phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoic acid;

ethyl 3-[[2-[[[3-[[amino(cyanoimino)methyl]amino]-phenyl]carbonyl]amino]acetyl]amino]-4-pentynoate;

- 921 -

(±) thyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-bis(trifluoromethyl)benzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(4,5-dihydro-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) 3,5-dibromo- $\beta$ -[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

- 922 -

(±) 3-bromo-5-chloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) [2-[2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]-carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) [2-[2-(2-hydroxyethoxy)-ethoxy]ethoxy]ethyl] 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3-bromo-5-chloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

ethyl (±) 3-bromo-5-chloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydropyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) 3,5-dichloro-β-[[2-[[[3-[(4,5-dihydro-1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]amino]acetyl]amino]-2-hydroxybenzenepropanoic acid;

- 923 -

(±) 3-bromo-5-chloro-β-[[2-[[[3-[(4,5-dihydro-1H-pyrrolidin-2-yl)amino]phenyl]carbonyl]amino]-acetyl]amino]-2-hydroxybenzenepropanoic acid;

β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

ethyl β-[[2-[[[3-[[5-phenyl-1H-imidazol-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate;

(±) ethyl 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3,5-dichloro-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino]phenyl]carbonyl]-amino]acetyl]amino]benzenepropanoic acid;

ethyl (±) 3,5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoate;

(±) 3-bromo-5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

ethyl(±) 3-bromo-5-dichloro-2-hydroxy-β-[[2-[[[3-[(1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]benzenepropanoate;

- 924 -

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-1,3-benzodioxole-5-propanoic acid;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-benzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoate;

$\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

(±) ethyl  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoate;

(±)  $\beta$ -[[2-[[[4-chloro-3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]pyridine-3-propanoic acid;

- 925 -

(±)  $\beta$ -[[2-[[[3,5-bis[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]pyridine-3-propanoic acid;

$\beta$ S-[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-4-pentynoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]-amino]-3,5-dichlorobenzenepropanoate;

(±)  $\beta$ -[[2-[[[3-[(3,4-dihydro-2H-pyrrol-5-yl)amino]phenyl]carbonyl]amino]acetyl]amino]-3,5-dichlorobenzenepropanoic acid;

(±) ethyl 3,5-dichloro- $\beta$ -[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoate;

(±) 3,5-dichloro- $\beta$ -[[2-[[[3-[(2,3,4,5-tetrahydropyridin-6-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

(±) 3,5-dichloro-2-hydroxy- $\beta$ -[[2-[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)amino]phenyl]carbonyl]amino]acetyl]amino]benzenepropanoic acid;

- 926 -

ethyl(±) 3,5-dichloro-2-hydroxy-β-[[2-  
[[[3-[(3,4,5,6-tetrahydro-2H-azepin-7-yl)-  
amino]phenyl]carbonyl]amino]acetyl]amino]-  
benzenepropanoate;

(±) ethyl β-[[2-[[[3-[(iminophenylmethyl)-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

βS-[[2-[[[3-[[imino(1-pyrrolidinyl)methyl]-  
amino]phenyl]carbonyl]amino]acetyl]-  
amino]-4-pentynoic acid;

(±) β-[[2-[[[3-[(iminophenylmethyl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

3,5-dichloro-β-[[2-[[[3-[[imino(1-  
piperidinyl)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoic acid;

ethyl 3,5-dichloro-β-[[2-[[[3-[[imino(1-  
piperidinyl)methyl]amino]phenyl]carbonyl]-  
amino]acetyl]amino]benzenepropanoate;

ethyl β-[[2-[[[3-[(4,5-dihydrothiazol-2-  
yl)amino]phenyl]carbonyl]amino]acetyl]-  
amino]pyridine-3-propanoate;

β-[[2-[[[3-[(4,5-dihydrothiazol-2-yl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

β-[[2-[[[3-[[benzoxazol-2-yl)amino]-  
phenyl]carbonyl]amino]acetyl]amino]-  
pyridine-3-propanoic acid;

- 927 -

ethyl  $\beta$ -[[2-[[[3-[[benzoxazol-2-yl)amino]-phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate.

$\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoic acid;

and

ethyl  $\beta$ -[[2-[[[3-[(5,6-dihydro-4H-thiazin-2-yl)-amino]phenyl]carbonyl]amino]acetyl]amino]-pyridine-3-propanoate.

43. A method according to Claim 41 wherein the condition treated is tumor metastasis.
44. A method according to Claim 42 wherein the condition treated is tumor metastasis.
45. A method according to Claim 41 wherein the condition treated is solid tumor growth.
46. A method according to Claim 42 wherein the condition treated is solid tumor growth.
47. A method according to Claim 41 wherein the condition treated is angiogenesis.
48. A method according to Claim 42 wherein the condition treated is angiogenesis.
49. A method according to Claim 41 wherein the condition treated is osteoporosis.
50. A method according to Claim 42 wherein the condition treated is osteoporosis.



- 928 -

51. A method according to Claim 41 wherein the condition treated is humoral hypercalcemia of malignancy.
52. A method according to Claim 42 wherein the condition treated is humoral hypercalcemia of malignancy.
53. A method according to Claim 41 wherein the condition treated is smooth muscle cell migration.
54. A method according to Claim 42 wherein the condition treated is smooth muscle cell migration.
55. A method according to Claim 53 wherein restenosis is inhibited.
56. A method according to Claim 54 wherein restenosis is inhibited.
57. A method according to Claim 41 wherein the condition treated is rheumatoid arthritis.
58. A method according to Claim 42 wherein the condition treated is rheumatoid arthritis.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/13500

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D213/55 A61K31/44 C07C279/18 A61K31/155 A61K31/36  
C07D405/10 A61K31/395 C07D223/12 C07D401/14 C07D207/16  
C07C275/28 A61K31/17 C07D401/10 C07D317/30

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 445 796 A (F.HOFFMANN-LA ROCHE) 11 September 1991 see page 7, line 37 - line 40; claim 1 ---	1-58
E	WO 96 26190 A (SMITHKLINE BEECHAM) 29 August 1996 see claim 1 ---	1-58
P,A	WO 96 00574 A (SMITHKLINE BEECHAM) 11 January 1996 see claim 1 ---	1-58
A	EP 0 643 072 A (TAKEDA) 15 March 1995 see claim 1 ---	1-58
A	WO 94 18981 A (MERCK & CO.) 1 September 1994 see page 28, line 22; claim 1 -----	1-58

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 December 1996

Date of mailing of the international search report

23.01.1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Gettins, M

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/13500

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-445796	11-09-91	CA-A- 2037153 IL-A- 97401 JP-B- 2501252 JP-A- 4217652 US-A- 5430024 US-A- 5273982 HR-A- 930353	10-09-91 15-03-95 29-05-96 07-08-92 04-07-95 28-12-93 30-06-96
WO-A-9626190	29-08-96	NONE	
WO-A-9600574	11-01-96	AU-A- 3001095 WO-A- 9600730	25-01-96 11-01-96
EP-A-643072	15-03-95	AU-A- 6477194 CA-A- 2126026 CN-A- 1098409 FI-A- 942881 HU-A- 70045 JP-A- 7157472 NO-A- 942274 US-A- 5550131	22-12-94 18-12-94 08-02-95 18-12-94 28-09-95 20-06-95 19-12-94 27-08-96
WO-A-9418981	01-09-94	AU-A- 6246594 BG-A- 99863 CA-A- 2155123 CN-A- 1118139 CZ-A- 9502108 EP-A- 0684823 FI-A- 953916 HU-A- 71796 JP-T- 8507072 NO-A- 953270 PL-A- 310386	14-09-94 29-02-96 01-09-94 06-03-96 14-02-96 06-12-95 21-08-95 28-02-96 30-07-96 19-10-95 11-12-95